

Table S1. Primers used for the detection of transcripts.

Target	Forward (5'-3')	Reverse (5'-3')
Human <i>RPS20</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>COL23A1</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>IL1R1</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>IL32</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>TLR4</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>CFH</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>C3</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>S100A9</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>IRF1</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>ADAM23</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>TNC</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
Human <i>SPINK5</i>	QuantiTect Primers (Qiagen, Hilden, Germany)	
HSV-1 <i>US12</i>	GTCCTCACGCCCCCTTTTAT	GAAATGGCGGACACCTTCCT
HSV-1 <i>UL19</i>	TCTCAGTCACAAAGCGGTCC	TTCAAGATCAGTCCCGTGGC
HSV-1 <i>UL23</i>	GGTCATGCTGCCCATAAGGT	CTCACCTCATCTTCGACCG
HSV-1 <i>UL27</i>	CACAGGGTCAGCTCGTGATT	CATACAGCGCCATGTCAACG
HSV-1 <i>UL42</i>	ACTCGCTTCTGGTTATGGGC	ACCGACTGAATTGCGAGTGT
HSV-1 <i>UL54</i>	CTCCAGTGCTACCTGAAGGC	TCCTTAATGTCCGCCAGACG




Figure S1: *COL23A1* overexpression enhances HSV-1 transcript levels. (A-F) Control and *COL23A1* overexpressing HaCaT cells were infected with 5 pfu/cell of HSV-1. *UL19*, *UL27*, *UL23*, *UL42*, *UL54* and *US12* mRNA levels were measured at 2 hpi, 4 hpi, 6 hpi, and 8 hpi by RT-qPCR and normalized to the housekeeping gene *RPS20* (black bars, mean $\pm$ standard deviation [SD]; n = 4 independently repeated experiments). Ctrl= HaCaT cells stably transduced with control lentiviral particles; COL23A1= HaCaT cells stably transduced with *COL23A1* lentiviral particles.