Supplementary 2., Fig. 1. Heatmap of genome-wide genetic correlations across 15 immunemediated diseases. A. Numbers in the cells show genetic correlations estimated using LD score regression method. Blue and red indicate positive and negative correlations respectively. B. Green and purple indicate positive and negative correlations respectively, asterisk means significant correlation between phenotypes (Bonferroni correction to number of comparisons). A.



Supplementary 2., Fig. 2. Genetic overlap beyond genetic correlations.

Fraction of disease-influencing genetic variants shared between two diseases with respect to the less polygenic disease in the pair estimated with Bivariate Gaussian causal mixture model (MiXeR) for 15 immune-mediated diseases.



Supplementary 2., Fig. 3. Local genetic correlations across immune-mediated diseases. Heatmap of local genetic correlations estimated with LAVA. The color indicates average correlation across all positively (top right) and negatively (bottom left) correlated regions. Numbers in cells indicate the number of regions with significant local generic correlation. The absence of number indicates that there were no regions with significant correlation.



Supplementary 2., Fig. 4. Genes shared between five or more immune-mediated diseases. For each disease, associated genes were identified using MAGMA.



Supplementary 2., Fig. 5. Tissue specificity analysis. For each disease, genes identified in MAGMA analysis were tested for tissue specific expression using gene expression data from 54 tissue types available in GTEx v8. Significant results are highlighted in red color. Analysis was performed using FUMA GENE2FUNC.



















CeD







MG







JIA





T1D







PSC



MS

Supplementary 2., Fig. 6. MAGMA-identified genes uniquely associated with diseases involved in factor 4 and linked to monogenic immune disorders and blood syndrome. Janus kinase 2 is involved in different blood disorders, *KLHDC8B* is involved in Hodgkin lymphoma, *RTEL1* in inborn error failure. *IRF1* and *CARD9* engage in different types of immunodeficiencies [Rosain et al., 2023, Wang et al., 2019, Salipante et al., 2009]. The *NOD2* gene plays a role in Blau syndrome, a rare autosomal dominant autoinflammatory syndrome classified as an autoinflammatory phenotype according to Tangye et al., 2022. Also, *STAT5B* gene is the monogenic cause of autosomal dominant growth hormone insensitivity syndrome with immune dysregulation-2.

