

Table S3. Participants' sociodemographic characteristics (n=78)

Characteristics	Ambulatory		On stretcher	
	EP	PT + EP	EP	PT + EP
Number of participants, <i>n</i> (%)	28 (35.9)	34 (43.6)	10 (12.8)	6 (7.7)
Age (yr), mean (SD)	39.2 (16.0)	35.0 (16.8)	58.0 (12.6)	45.3 (18.8)
Sex, <i>n</i> females (%)	8 (28.6)	18 (52.9)	4 (40.0)	4 (66.7)
Triage category* in ED, <i>n</i> (%)				
Urgent (P3)	8 (28.6)	10 (29.4)	8 (80.0)	6 (100.0)
Semi urgent (P4)	19 (67.9)	24 (70.6)	2 (20.0)	0 (0.0)
Non urgent (P5)	1 (3.6)	0 (0.0)	0 (0.0)	0 (0.0)
ED length of stay (h), mean (SD)	6.4 (2.9)	6.2 (3.9)	9.2 (8.3)	9.5 (7.0)
Other health condition, yes (%)	13 (46.4)	21 (61.8)	10 (100.0)	5 (83.3)

EP: emergency physician; PT: physiotherapist; yr: year; SD: standard deviation; ED: emergency department; h: hour

* According to the Canadian Triage and Acuity Scale (CTAS)

More information on the participants included in this study can be found in the following article:

Gagnon R, Perreault K, Berthelot S, et al. Direct-access physiotherapy to help manage patients with musculoskeletal disorders in an emergency department: Results of a randomized controlled trial. Acad Emerg Med 2021;28(8):848-858. DOI: 10.1111/acem.14237.