Table S4. Average cost of an emergency department visit by care model and orientation

Costs (in 2019 Canadian dollars) (n=78) Ambulatory (n=62) Stretcher (n=16) PT + EPEP PT + EPEP n = 28n = 34n=10n=6Absolute difference Absolute difference Costs\* - Median time values Costs\* - Median time values Care processes p p (95% CI) (95% CI) 145.67 214.19 559.91 642.68 Average cost of 68.52 82.77 <.01 .63 (354.58, 828.67) the ED visit (117.20, 174.97) (190.84, 238.22) (29.80, 104.36) (389.93, 996.39)(-303.11, 492.83)336.49 98.07 179.80 81.73 203.12 133.37 ED check-in<sup>¶</sup> <.01 <.01 (80.31, 117.73) (166.30, 192.69)(58.81, 103.84) (170.25, 254.27) (232.99, 477.36)(16.64, 280.94) 235.57 243.80 0.00 0.008.23 ED stay<sup>●</sup> .93 (---) (---) (124.67, 377.55)(116.62, 391.63)(-198.28, 177.55)25.94 13.99 11.95 68.48 34.06 34.42 **Imaging** .30 .97 (0.00, 92.54)(15.86, 37.33)(10.07, 18.11)(0.97, 23.86)(26.84, 114.10)(-98.59, 36.14)2.92 4.53 1.61 16.63 10.81 5.82 Laboratory tests .99 .98 (0.00, 6.66)(0.00, 14.00)(-5.59, 11.47)(0.00, 55.42)(0.00, 21.20)(-44.29, 18.27)0.99 1.26 0.27 18.17 9.11 9.06 Medication .98 .31 (0.33, 1.81)(0.11, 2.76)(-1.26, 1.96)(9.59, 28.99)(3.14, 15-56)(-21.14, 1.61)

.99

17.94

(0.00, 59.80)

8.40

(0.00, 18.77)

9.54

(-54.49, 15.41)

.99

EP: emergency physician; PT: physiotherapist; CI: confidence interval; ED: emergency department

14.61

(7.90, 22.02)

Musculoskeletal

disorders†

17.75

(8.14, 29.97)

3.14

(-17.47, 8.29)

<sup>\*</sup>Costs are presented as mean (95% confidence intervals obtained via non-parametric Bootstrap).

<sup>¶</sup> Includes ED arrival (clerk, nurse, nursing assistant), triage reassessment (if needed; nurse), assessment (emergency physician; if needed: physiotherapist, internist, social worker, occupational therapist), reassessment (if needed; emergency physician, physiotherapist), point-of-care ultrasound (if needed; emergency physician).

<sup>•</sup> Since all the parameters needed to calculate the ED stay cost are fixed (hours billed, patient ratio, ED length of stay), the average costs obtained are the same for all three scenarios.

<sup>&</sup>lt;sup>†</sup> Musculoskeletal disorders care processes include supplying and advising on use for walking aids and orthoses (crutches, cane, walker or walking boot), and splinting (plaster cast, Jones splint, thoracobrachial splint, ankle splint or Zimmer splint).

Table S5. Average cost of an emergency department visit by care model and orientation according to different care process durations (sensitivity analyses)

	Costs (in 2019 Canadian dollars) (n=78)					
	EP	PT + EP		EP	PT + EP	
Care processes	n=28	n=34		n=28	n=34	
Ambulatory (n=62)	Costs* – Q1 time values		p	Costs* – Q3 time values		p
Average cost of the	103.13	170.61	<.01	203.51	273.03	.02
ED visit	(82.49, 123.81)	(153.97, 187.72)	<b>\.</b> 01	(162.97, 248.01)	(240.48, 305.58)	.02
ED check-in¶	61.47	141.60	<.01	148.09	232.10	<.01
	(51.52, 72.80)	(133.94, 148.73)		(119.21, 179.89)	(210.04, 253.38)	
ED stay•	0.00	0.00		0.00	0.00	
	()	()		()	()	
Imaging	22.67	12.24	.30	29.90	15.51	.27
	(13.86, 32.69)	(8.95, 15.62)		(18.55, 43.13)	(11.18, 20.05)	
Laboratory tests	2.58	4.30	.99	3.54	4.88	.99
	(0.00, 5.91)	(0.00, 13.30)		(0.00, 7.95)	(0.00, 15.09)	
Medication	0.66	0.82	.97	1.41	1.82	.97
	(0.23, 1.19)	(0.07, 1.77)	.,,	(0.47, 2.60)	(0.15, 3.98)	.,,
Musculoskeletal	15.74	11.65	.99	20.57	18.73	.99
disorders <sup>†</sup>	(6.70, 28.01)	(5.95, 17.94)	.,,	(10.07, 33.83)	(10.64, 27.90)	•,,,
	EP	PT + EP		EP	PT + EP	
	n=10	n=6		n=10	n=6	
Stretcher (n=16)	Costs* – Q1 time values		р	Costs* – Q3 time values		p
Average cost of the	483.23	582.82	-	653.53	720.67	-
ED visit	(287.64, 739.18)	(347.21, 926.14)	.55	(434.60, 936.88)	(442.07, 1,094.60)	.73
ED check-in¶	143.47	285.45	<.01	278.60	400.42	.02
	(115.75, 189.12)	(197.66, 415.20)		(239.92, 336.03)	(276.67-549.75)	
ED stay•						
Imaging	58.98	30.49	07	76.78	40.07	.97
	50.70	30.77				
mugmg	(22.51, 99.59)	(0.00, 83.48)	.97	(30.26, 127.75)	(0.00, 108.94)	.97
				(30.26, 127.75) 18.36	(0.00, 108.94) 13.71	
	(22.51, 99.59)	(0.00, 83.48)	.97			.98
Laboratory tests	(22.51, 99.59) 15.68	(0.00, 83.48) 9.23	.98	18.36	13.71	.98
Laboratory tests	(22.51, 99.59) 15.68 (0.00, 52.25)	(0.00, 83.48) 9.23 (0.00, 18.45)		18.36 (0.00, 61.21)	13.71 (0.00, 26.41)	
Laboratory tests  Medication  Musculoskeletal	(22.51, 99.59) 15.68 (0.00, 52.25) 12.99	(0.00, 83.48) 9.23 (0.00, 18.45) 5.91	.98 .24	18.36 (0.00, 61.21) 24.75	13.71 (0.00, 26.41) 13.12	.98 .35
Laboratory tests Medication	(22.51, 99.59) 15.68 (0.00, 52.25) 12.99 (6.26, 21.82)	(0.00, 83.48) 9.23 (0.00, 18.45) 5.91 (2.06, 9.92)	.98	18.36 (0.00, 61.21) 24.75 (13.65, 38.39)	13.71 (0.00, 26.41) 13.12 (4.51, 22.43)	.98

EP: emergency physician; PT: physiotherapist; CI: confidence interval; ED: emergency department; Q1: first quartile; Q3: third quartile

<sup>\*</sup>Costs are presented as mean (95% confidence intervals obtained via non-parametric Bootstrap).

<sup>¶</sup> Includes ED arrival (clerk, nurse, nursing assistant), triage reassessment (if needed; nurse), assessment (emergency physician; if needed: physiotherapist, internist, social worker, occupational therapist), reassessment (if needed; emergency physician, physiotherapist), point-of-care ultrasound (if needed; emergency physician).

• Since all the parameters needed to calculate the ED stay cost are fixed (hours billed, patient ratio, ED length of stay), the average costs obtained are the same for all three scenarios.

<sup>†</sup> Musculoskeletal disorders care processes include supplying and advising on use for walking aids and orthoses (crutches, cane, walker or walking boot), and splinting (plaster cast, Jones splint, thoracobrachial splint, ankle splint or Zimmer splint).