

1 **Exploring and mitigating potential bias when genetic instrumental variables are associated with**
2 **multiple non-exposure traits in Mendelian randomization**

3 Qian Yang, Eleanor Sanderson, Kate Tilling, M Carolina Borges, Deborah A Lawlor

4

5 **SUPPLEMENTARY MATERIAL**

6 Supplementary Methods

7 Supplementary Figure 1. The number of participants in one- and two-sample Mendelian
8 randomization of our real data example

9 Supplementary Figure 2. Distributions of polygenetic risk score for insomnia across 22 UK Biobank
10 assessment centres

11 Supplementary Figure 3. Scatter plots of two-sample Mendelian randomization for the effect of
12 insomnia on birthweight using MR-Base

13 Supplementary Figure 4. Scatter plots of two-sample Mendelian randomization for the effect of
14 insomnia on birthweight using MR-TRYX

15 Supplementary Table 1. The UK Biobank data fields of variables used in this study

16 Supplementary Table 2. Results for age at first live birth adjusting for genetic array, participants' age
17 and birthplace, and UK Biobank assessment centres

18 Supplementary Table 3. Strengths of polygenetic risk score (PRS)

19 Supplementary Data 1. Information of genetic variants used as instrumental variables in real data
20 example

21 Supplementary Data 2. Original output from 'some invalid some Valid IV Estimator' for the effect of
22 insomnia on birthweight

23 Supplementary Data 3. Genome-wide significant associations of insomnia-related variants extracted
24 from Phenoscanner

25 *Supplementary Data are in a separate file.

26 **Supplementary Methods**

27 ***Genetic instrumental variables (IVs) for insomnia***

28 We used single nucleotide polymorphisms (SNPs) identified by the largest genome-wide association
29 study (GWAS) combining UK Biobank (UKB) with 23andMe participants⁽¹⁾ due to its sample size
30 (N=709,986 women), provision of female-specific results and a low proportion (29%) of overlapped
31 participants with UKB women to try to avoid winner's curse.⁽²⁾ This GWAS reported 83 female-
32 specific SNPs robustly associated with insomnia (P-value < 5×10^{-8}). From these, we removed 3 SNPs
33 that were correlated to other variants using MR-Base 'clumping' function ($R^2=0.01$, referring to the
34 European samples from the 1000 genomes project).⁽³⁾ We extracted genotypes from UKB for the
35 remaining 80 SNPs and derived an unweighted polygenic risk score (PRS), by adding up the number
36 of insomnia risk-raising alleles. We used the PRS as the IV in one-sample Mendelian randomization
37 (MR).

38 ***Genetic IVs for height, body mass index (BMI), age at first live birth, education, frequency of*** 39 ***alcohol intake and ever smoking***

40 We aimed to use GWAS conducted in only women of European descent which did not overlap with
41 UKB. We searched for genome-wide significant (P-value < 5×10^{-8}) SNPs and performed 'clumping'
42 ($R^2=0.01$, referring to the European samples from the 1000 genomes project) for those traits on MR-
43 Base, if their GWAS were included. The Genetic Investigation of Anthropometric Traits consortium
44 identified 54 and 38 SNPs associated with female height (N=147,746, dataset ID '97' on MR-Base)⁽⁴⁾
45 and female BMI (N=171,977, dataset ID '974' on MR-Base),⁽⁵⁾ respectively. Social Science Genetic
46 Association Consortium identified 6 and 19 SNPs associated with female age at first live birth
47 (N=154,839, not on MR-Base)⁽⁶⁾ and female year of schooling (N=182,286, dataset ID '1011' on MR-
48 Base),⁽⁷⁾ respectively. These four GWAS were conducted in non-UKB settings. Neale Lab's GWAS
49 identified 44 and 40 SNPs associated with frequency of alcohol intake (N=336,965, dataset ID 'UKB-
50 a:25' on MR-Base) and ever smoking (N=336,067, dataset ID 'UKB-a:236' on MR-Base) in UKB men
51 and women.⁽⁸⁾ We derived weighted PRS as IVs for one-sample MR using effect sizes reported by the
52 corresponding GWAS as the weights. To minimize bias from internal weight,⁽⁹⁾ we also followed
53 previous MR studies^(10, 11) to repeat our analyses using rs1229984 (*ADH1B*) and rs698 (*ADH1C*) for
54 frequency of alcohol intake and rs6265 (*BDNF*) for ever smoking, which were identified in non-UKB
55 settings.^(12, 13)

56 ***Exploring the role of population stratification***

57 Individual level data allow us to check for population stratification. We tested associations of
58 insomnia PRS with maternal age at recruitment and birthplace (including longitude and latitude) and
59 compared means of PRS across 22 UKB study centres using ANOVA. We also compared each non-
60 exposure trait association according to PRS using (i) a crude model, (ii) a model adjusting for genetic
61 array and top 40 principal components (PCs), (iii) a model further adjusting for participants' age at
62 recruitment and birthplace, and (iv) a model adjusting for all these covariates plus UKB study
63 centres. We obtained differences in mean non-exposure traits per allele increase in PRS from a linear
64 regression, except for ever smoking where we applied logistic regression.

65 ***Univariable MR to assess bias due to horizontal pleiotropy***

66 We used univariable MR to explore (i) the associations of the six non-exposure traits (W) with
67 birthweight (Y), (ii) the causal directions between insomnia (X) and W, and (3) the association of X
68 with Y for comparison. In one-sample setting, we applied two-stage least squares (TSLS).⁽¹⁴⁾ In the
69 first stage, the exposure (a continuous or binary variable, x_i) was regressed on its PRS (z_i) in a linear
70 model (equation 1). In the second stage, the outcome (a continuous or binary, y_i) was regressed on

71 the fitted values for exposure (\hat{x}_i) from the first stage in a linear model (equation 2). ε_{x_i} and ε_{y_i}
72 represented the independent error terms. We used a linear model rather than logit model (which
73 would be more common with a binary exposure) to avoid non-collapsibility of odds ratio and difficult
74 interpretation of the units (e.g. per doubling of probability of binary exposure).⁽¹⁵⁾

75
$$x_i = \alpha_0 + \alpha_1 \times z_i + \varepsilon_{x_i} \text{ (equation 1)}$$

76
$$y_i = \beta_0 + \beta_1 \times \hat{x}_i + \varepsilon_{y_i} \text{ (equation 2)}$$

77 The unit of TSLS estimates correspond to difference in mean per one unit increase in the exposure.
78 Specifically, the unit of birthweight is gram. We standardized height, BMI and age at first live birth to
79 make their estimates comparable to each other, and 1 SD is equivalent to 6 cm in height, 5 kg/m² in
80 BMI and 5 years in age at first live birth. For education and frequency of alcohol intake, which are
81 ordered categorical, one-unit increase represents one level change from a lower category to a higher
82 category. For binary variables, one-unit increase in ever smoking represents comparing ever smokers
83 to never smokers, and one-unit increase in insomnia represents comparing participants reporting
84 that they “usually” experience insomnia to “sometimes/rarely/never”. We obtained differences in
85 mean (or absolute differences in risk of binary variables) together with 95% confidence intervals
86 using ‘ivreg’ package in R, which can give a correct standard error considering the uncertainty in
87 both regression stages.

88 We examined the strength of PRS via F-statistic in each first stage regression,⁽¹⁶⁾ with results shown
89 in Supplementary Table 3. For the effect of insomnia on birthweight, we assessed between SNP
90 heterogeneity via Sargan test which is an ‘overidentifying’ test and need individual SNPs to be IVs in
91 TSLS. Rejection of its null hypothesis suggests at least one invalid IV.⁽¹⁷⁾

92 Since the GWAS of insomnia included UKB,⁽¹⁾ we followed a previous study that had the same
93 problem⁽¹⁸⁾ to conduct our two-sample MR. We randomly split our UKB women (N=208,171) into
94 two sets (N_A=104,041 and N_B=104,130, see Supplementary Figure 1). In each dataset, we obtained
95 SNP-specific associations with each trait (i.e. IV_X-X, IV_X-Y, IV_X-W, IV_W-X, IV_W-Y and IV_W-W) by running
96 linear regressions. For W-Y associations, we used IV_W-W from dataset A and IV_W-Y from dataset B (A
97 on B) and vice versa (B on A) in inverse variance weighted (IVW) analyses. Similarly, for W-X
98 associations, we used IV_W-W from dataset A and IV_W-X from dataset B (A on B) and vice versa (B on
99 A) in IVW; for X-W associations, we used IV_X-X from dataset A and IV_X-W from dataset B (A on B) and
100 vice versa (B on A) in IVW; for the X-Y association, we used IV_X-X from dataset A and IV_X-Y from
101 dataset B (A on B) and vice versa (B on A) in IVW. We used ‘TwoSampleMR’ package in R to conduct
102 the IVW analyses. At last, we pooled the MR estimates from the two together for each association.
103 For the effect of X on Y, we assessed between SNP heterogeneity via Cochran’s Q statistic via MR-
104 Base.

105 In one- and two-sample MR, we included genetic array and top 40 PCs as covariates, given further
106 adjustments for participants’ age, birthplace and study centre showed relatively similar associations
107 in Figure 1. We conducted sensitivity analyses for age at first live birth to include those as covariates,
108 with results shown in Supplementary Table 2.

109 **Multivariable MR to account for horizontal pleiotropy**

110 We would conduct multivariable MR of effects of (i) insomnia and age at first live birth, (ii) insomnia
111 and education, (iii) insomnia and ever smoking, and (iv) insomnia, age at first live birth, education
112 and ever smoking on birthweight. In one-sample setting, we used PRS for X and W as their IVs in
113 TSLS, and their strengths were assessed via F-statistics (results shown in Supplementary Table 2).⁽¹⁹⁾

114 In two-sample setting, we conducted multivariable MR using IV_{X-X} , IV_{X-W} , IV_{W-W} , IV_{W-X} from dataset
115 A and IV_{X-Y} , IV_{W-Y} from dataset B (A on B) and vice versa (B on A). Finally, we pooled the MR
116 estimates from the two together for each insomnia-birthweight association.


117 ***Sensitivity analyses***

118 In one-sample MR, we applied some invalid some valid instrumental variable estimator (sisVIVE) by
119 using the 'sisVIVE' package in R.⁽²⁰⁾ The package requires our data in a matrix format with 3 columns
120 (i.e. PRS, insomnia, birthweight) \times 165,184 rows (i.e. the number of participants). sisVIVE can identify
121 one invalid IV at a time and provide a difference in mean birthweight with correction for that. Since
122 we have 80 variants for insomnia, sisVIVE provided 80 differences in mean birthweight with
123 correction for 0, 1, 2 78 and 79 invalid IVs (full results in Supplementary Data 2).


124 In two-sample MR, we used 'TwoSampleMR' package in R⁽³⁾ to obtain estimates from MR-Egger,⁽²¹⁾
125 weighted median⁽²²⁾ and weighted mode⁽²³⁾ approaches, 'MRPRESSO' package in R to obtain
126 estimates from MR Pleiotropy RESidual Sum and Outlier method,⁽²⁴⁾ and 'tryx' package in R to obtain
127 estimates from MR Treasure Your eXceptions.⁽²⁵⁾ In each method, we used IV_{X-X} from dataset A and
128 IV_{X-Y} from dataset B (A on B) and vice versa (B on A), and finally meta-analysed the MR estimates
129 from the two.

130

131 **Supplementary Figure 1. The number of participants in one- and two-sample Mendelian randomization of our real data example**



132




134

135 Boxplot shows median, interquartile range (IQR), minimum (25th percentile – 1.5*IQR), maximum (75th percentile + 1.5*IQR) and outliers.


136 **Supplementary Figure 3. Scatter plots of two-sample Mendelian randomization for the effect of insomnia on birthweight using MR-Base**

137 (a) Dataset A on dataset B (IVW Q statistic = 101, P = 0.048; MR-Egger intercept = 0.275, SE = 0.797, P = 0.732)



138

139 (b) Dataset B on dataset A (IVW Q statistic = 157, $P = 3.85 \times 10^{-7}$; MR-Egger intercept = -0.315, SE = 1.041, $P = 0.763$)




140

141


142 **Supplementary Figure 4. Scatter plots of two-sample Mendelian randomization for the effect of insomnia on birthweight using MR-TRYX**

143 (a) Dataset A on dataset B (Q statistic = 84 in the two outliers removed models and Q = 91 in the outlier adjusted model)



144

145 (b) Dataset B on dataset A (Q = 117 in the two outliers removed models and Q = 141 in the outlier adjusted model)



146

147

148 **Supplementary Table 1. The UK Biobank data fields of variables used in this study**

Variable	Field ID	Code in this study	N
Exposure			
Maternal insomnia	1200	"Do you have trouble falling asleep at night or do you wake up in the middle of the night?", with answers "never/rarely"=0, "sometimes"=0 and "usually"=1.	207,924
Outcome			
Offspring birthweight	2744	"What was the birth weight of your first child in pounds? (do not include twins)", with answers ranging from 2 to 15. We converted its unit to grams.	165,254
Maternal characteristics			
Height	50	A continuous variable, mean≈163 cm, SD≈6 cm	207,704
Body mass index	21001	A continuous variable, mean≈27 kg/m ² , SD≈5 kg/m ²	207,477
Age at first live birth	2754	"How old were you when you had your FIRST child?" A continuous variable, mean≈25 years, SD≈5 years	140,503
Education	6138	"Which of the following qualifications do you have? (You can select more than one)", with answers "College or University degree"=3, "A levels/AS levels or equivalent"=2, "O levels/GCSEs or equivalent"=1, "CSEs or equivalent"=1, "NVQ or HND or HNC or equivalent"=1, "Other professional qualifications eg: nursing, teaching"=2, "None of the above"=1.	206,055
Frequency of alcohol intake	1558	"About how often do you drink alcohol?", with answers "Daily or almost daily"=6, "Three or four times a week"=5, "Once or twice a week"=4, "One to three times a month"=3, "Special occasions only"=2, "Never"=1.	207,872
Ever smoking	20116	"Never" = 0, "Previous" = 1, "Current" = 1.	207,281
Covariates			
Assessment centre	54	A categorical variable	208,171
Age	21003	A continuous variable	208,171
Place of birth in UK	129, 130	Two continuous variables (longitude and latitude)	189,470

149 We coded "Prefer not to answer" as missing. All phenotypic values in the analyses were no less than 0. Details of how these variables were assessed can be
150 found on <http://biobank.ctsu.ox.ac.uk/crystal/search.cgi>.

151 **Supplementary Table 2. Results for age at first live birth adjusting for genetic array, participants' age and birthplace, and UK Biobank assessment centres**

Mendelian randomization	Setting	Model 1	Model 2
Univariable			
Age at first live birth (SD)→ birthweight (grams)	1	85.2 (-11.0, 181.5)	96.1 (-5.1, 197.3)
	2	87.4 (-9.5, 184.4)	91.2 (-7.7, 190.0)
Age at first live birth (SD)→ insomnia	1	-0.084 (-0.166, -0.002)	-0.077 (-0.163, 0.009)
	2	-0.123 (-0.197, -0.056)	-0.148 (-0.222, -0.073)
Insomnia→ age at first live birth (SD)	1	-1.025 (-1.213, -0.838)	-0.991 (-1.178, -0.804)
	2	-0.826 (-1.048, -0.605)	-0.769 (-0.982, -0.555)
Multivariable			
Insomnia + age at first live birth→ birthweight (grams)	1	18.0 (-131.3, 167.2)	23.1 (-127.6, 173.8)
	2	-111.9 (-255.7, 32.0)	-99.4 (-244.1, 45.4)
Insomnia + age at first live birth + education + being ever smokers→ birthweight (grams)	1	-37.5 (-224.4, 149.4)	-56.5 (-243.0, 130.0)
	2	-52.0 (-155.2, 51.2)	-48.7 (-152.9, 55.4)

152 Estimates are differences in mean outcome per unit increase in exposure. One unit of age at first live birth is 1 SD (5 years) in univariable Mendelian
 153 randomization and 1 year in multivariable Mendelian randomization. Education has 3 levels; being ever smokers and insomnia are binary. Model 1 adjusted
 154 for genetic array and top 40 principal components (presented in Figure 2), while Model 2 further adjusted for participants' age, birthplace and assessment
 155 centre.

156

157 **Supplementary Table 3. Strengths of polygenetic risk score (PRS)**

Mendelian randomization	F-statistics ^a	
Univariable		
PRS of 54 variants → height → birthweight	9376	
PRS of 38 variants → body mass index → birthweight	1,946	
PRS of 6 variants → age at first live birth → birthweight	147	
PRS of 19 variants → education → birthweight	574	
PRS of 44 variants → alcohol consumption frequency → birthweight	1,264	
PRS of 2 variants → alcohol consumption frequency → birthweight	181 ^b	
PRS of 40 variants → being ever smokers → birthweight	604	
rs6265 (<i>BDNF</i>) → being ever smokers → birthweight	8 ^c	
PRS of 54 variants → height → insomnia	11,886	
PRS of 38 variants → body mass index → insomnia	2,668	
PRS of 6 variants → age at first live birth → insomnia	149	
PRS of 19 variants → education → insomnia	744	
PRS of 44 variants → alcohol consumption frequency → insomnia	1,590	
PRS of 2 variants → alcohol consumption frequency → insomnia	212 ^b	
PRS of 40 variants → being ever smokers → insomnia	777	
rs6265 (<i>BDNF</i>) → being ever smokers → insomnia	7 ^c	
PRS of 80 variants → insomnia → height	686	
PRS of 80 variants → insomnia → body mass index	682	
PRS of 80 variants → insomnia → age at first live birth	494	
PRS of 80 variants → insomnia → education	671	
PRS of 80 variants → insomnia → alcohol consumption frequency	690	
PRS of 80 variants → insomnia → being ever smokers	688	
PRS of 80 variants ^d → insomnia → birthweight	591	
Multivariable		
PRS ₁ + PRS ₂ → insomnia + age at first live birth → birthweight		
PRS ₁ of 80 variants for insomnia	245	73
PRS ₂ of 6 variants for age at first live birth	132	64
PRS ₁ + PRS ₃ → insomnia + education → birthweight		
PRS ₁ of 80 variants for insomnia	289	286
PRS ₃ of 19 variants for education	299	368
PRS ₁ + PRS ₄ → insomnia + being ever smokers → birthweight		
PRS ₁ of 80 variants for insomnia	300	299
PRS ₄ of 40 variants for being ever smokers	320	333
PRS ₁ + PRS ₂ + PRS ₃ + PRS ₄ → insomnia + age at first live birth + education + being ever smokers → birthweight		
PRS ₁ of 80 variants for insomnia	123	7
PRS ₂ of 6 variants for age at first live birth	105	5
PRS ₃ of 19 variants for education	150	5
PRS ₄ of 40 variants for being ever smokers	127	25

158 ^a F-statistics for the same PRS may be slightly different due to different sample sizes in the analyses.

159 ^b Little evidence was found for a causal effect.

160 ^c No further analyses were conducted using this instrumental variable due to its small F-statistic.

161 ^d When 80 individual SNPs were used as instrumental variables in two-stage least squares, Sargan test suggests invalid IVs ($P = 3 \times 10^{-8}$).

163 **References**

- 164 1. Jansen PR, Watanabe K, Stringer S, *et al.* Genome-wide analysis of insomnia in 1,331,010
165 individuals identifies new risk loci and functional pathways. *Nat Genet.* 2019;**51**:394-403.
- 166 2. Lawlor DA. Commentary: Two-sample Mendelian randomization: opportunities and
167 challenges. *Int J Epidemiol.* 2016;**45**:908-915.
- 168 3. Hemani G, Zheng J, Elsworth B, *et al.* The MR-Base platform supports systematic causal
169 inference across the human phenome. *Elife.* 2018;**7**.
- 170 4. Randall JC, Winkler TW, Kutalik Z, *et al.* Sex-stratified genome-wide association studies
171 including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. *PLoS*
172 *Genet.* 2013;**9**:e1003500.
- 173 5. Locke AE, Kahali B, Berndt SI, *et al.* Genetic studies of body mass index yield new insights for
174 obesity biology. *Nature.* 2015;**518**:197-206.
- 175 6. Barban N, Jansen R, de Vlaming R, *et al.* Genome-wide analysis identifies 12 loci influencing
176 human reproductive behavior. *Nat Genet.* 2016;**48**:1462-1472.
- 177 7. Okbay A, Beauchamp JP, Fontana MA, *et al.* Genome-wide association study identifies 74 loci
178 associated with educational attainment. *Nature.* 2016;**533**:539-542.
- 179 8. Nealelab. We're thrilled to announce an updated GWAS analysis of the UK Biobank.
180 2018:<http://www.nealelab.is/uk-biobank>.
- 181 9. Hartwig FP, Davies NM. Why internal weights should be avoided (not only) in MR-Egger
182 regression. *Int J Epidemiol.* 2016;**45**:1676-1678.
- 183 10. Lawlor DA, Nordestgaard BG, Benn M, *et al.* Exploring causal associations between alcohol
184 and coronary heart disease risk factors: findings from a Mendelian randomization study in the
185 Copenhagen General Population Study. *Eur Heart J.* 2013;**34**:2519-2528.
- 186 11. Gibson M, Munafo MR, Taylor AE, *et al.* Evidence for Genetic Correlations and Bidirectional,
187 Causal Effects Between Smoking and Sleep Behaviors. *Nicotine Tob Res.* 2019;**21**:731-738.
- 188 12. Tolstrup JS, Nordestgaard BG, Rasmussen S, *et al.* Alcoholism and alcohol drinking habits
189 predicted from alcohol dehydrogenase genes. *Pharmacogenomics J.* 2008;**8**:220-227.
- 190 13. Genome-wide meta-analyses identify multiple loci associated with smoking behavior. *Nat*
191 *Genet.* 2010;**42**:441-447.
- 192 14. Burgess S, Dudbridge F, Thompson SG. Combining information on multiple instrumental
193 variables in Mendelian randomization: comparison of allele score and summarized data methods.
194 *Stat Med.* 2016;**35**:1880-1906.
- 195 15. Burgess S, Labrecque JA. Mendelian randomization with a binary exposure variable:
196 interpretation and presentation of causal estimates. *Eur J Epidemiol.* 2018;**33**:947-952.
- 197 16. Burgess S, Small DS, Thompson SG. A review of instrumental variable estimators for
198 Mendelian randomization. *Stat Methods Med Res.* 2017;**26**:2333-2355.
- 199 17. Burgess S, Scott RA, Timpson NJ, *et al.* Using published data in Mendelian randomization: a
200 blueprint for efficient identification of causal risk factors. *Eur J Epidemiol.* 2015;**30**:543-552.
- 201 18. Henry A, Katsoulis M, Masi S, *et al.* The relationship between sleep duration, cognition and
202 dementia: a Mendelian randomization study. *Int J Epidemiol.* 2019.
- 203 19. Sanderson E, Davey Smith G, Windmeijer F, *et al.* An examination of multivariable Mendelian
204 randomization in the single-sample and two-sample summary data settings. *Int J Epidemiol.* 2018.
- 205 20. Kang H, Zhang A, Cai TT, *et al.* Instrumental Variables Estimation With Some Invalid
206 Instruments and its Application to Mendelian Randomization. *Journal of the American Statistical*
207 *Association.* 2016;**111**:132-144. <https://doi.org/10.1080/01621459.2014.994705>.
- 208 21. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments:
209 effect estimation and bias detection through Egger regression. *Int J Epidemiol.* 2015;**44**:512-525.
- 210 22. Bowden J, Davey Smith G, Haycock PC, *et al.* Consistent Estimation in Mendelian
211 Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol.*
212 2016;**40**:304-314.

- 213 23. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data Mendelian
214 randomization via the zero modal pleiotropy assumption. *Int J Epidemiol.* 2017;**46**:1985-1998.
- 215 24. Verbanck M, Chen CY, Neale B, *et al.* Detection of widespread horizontal pleiotropy in causal
216 relationships inferred from Mendelian randomization between complex traits and diseases. *Nat*
217 *Genet.* 2018;**50**:693-698.
- 218 25. Cho Y, Haycock PC, Gaunt TR, *et al.* MR-TRYX: Exploiting horizontal pleiotropy to infer novel
219 causal pathways. *bioRxiv.* 2018:<https://doi.org/10.1101/476085>.

220