

Supplemental tables

Table A. Pairwise comparison of odds ratios for asthma after two years of age following early-life respiratory infection by racial/ethnic group.

Illness	Group	Group Pair		
		<i>African American</i>	<i>Combined</i>	<i>Mexican</i>
<i>URI</i>	<i>Combined</i>	0.760	-	-
	<i>Mexican</i>	0.067	0.066	-
	<i>Puerto Rican</i>	0.797	0.487	0.031
<i>Pneumonia</i>	<i>Combined</i>	0.428	-	-
	<i>Mexican</i>	0.882	0.578	-
	<i>Puerto Rican</i>	0.037	0.089	0.059
<i>Bronchitis</i>	<i>Combined</i>	0.081	-	-
	<i>Mexican</i>	0.356	0.382	-
	<i>Puerto Rican</i>	0.011	0.168	0.057
<i>Bronchiolitis/RSV</i>	<i>Combined</i>	0.435	-	-
	<i>Mexican</i>	0.733	0.147	-
	<i>Puerto Rican</i>	0.309	0.623	0.085
<i>Any Listed</i>	<i>Combined</i>	0.581	-	-
	<i>Mexican</i>	0.443	0.136	-
	<i>Puerto Rican</i>	0.125	0.179	0.019

5 **Table B.** Descriptive statistics of the study population including those diagnosed with asthma before age two (N = 3,722).

	Puerto Rican				Mexican American				African American				Combined			
	Case		Control		Case		Control		Case		Control		Case		Control	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Number of subjects	791	100	783	100	499	100	504	100	699	100	446	100	1989	100	1733	100
Males	439	55.5	361	46.1	288	57.7	210	41.7	370	52.9	187	41.9	1097	55.2	758	43.7
Underweight at birth	84	10.6	61	7.8	39	7.8	35	6.9	73	10.4	45	10.1	196	9.9	141	8.1
In-utero smoke exposure	53	6.7	40	5.1	17	3.4	9	1.8	139	19.9	54	12.1	209	10.5	103	5.9
Breastfed	428	54.1	432	55.2	379	76	401	79.6	399	57.1	244	54.7	1206	60.6	1077	62.1
Number of older siblings																
0	197	24.9	200	25.5	169	33.9	195	38.7	393	56.2	208	46.6	759	38.2	603	34.8
1	240	30.3	274	35	185	37.1	176	34.9	161	23	132	29.6	586	29.5	582	33.6
2 or more	354	44.8	309	39.5	145	29.1	133	26.4	145	20.7	106	23.8	644	32.4	548	31.6
Socioeconomic status*																
high	232	29.3	271	34.6	142	28.5	116	23	240	34.3	160	35.9	614	30.9	547	31.6
medium	142	18	136	17.4	104	20.8	84	16.7	183	26.2	137	30.7	429	21.6	357	20.6
low	417	52.7	376	48	253	50.7	304	60.3	276	39.5	149	33.4	946	47.6	829	47.8
Recruitment region																
Chicago	36	4.6	23	2.9	178	35.7	180	35.7	-	-	-	-	214	10.8	203	11.7
Houston	2	0.3	-	-	126	25.3	91	18.1	-	-	-	-	128	6.4	91	5.3
New York	64	8.1	34	4.3	28	5.6	64	12.7	-	-	-	-	92	4.6	98	5.7
San Francisco Bay Area	1	0.1	1	0.1	167	33.5	169	33.5	699	100	446	100	867	43.6	616	35.5
Puerto Rico	688	87	725	92.6	-	-	-	-	-	-	-	-	688	34.6	725	41.8
URI	254	32.1	35	4.5	55	11	27	5.4	200	28.6	16	3.6	509	25.6	78	4.5
Pneumonia	71	9	5	0.6	49	9.8	11	2.2	64	9.2	14	3.1	184	9.3	30	1.7
Bronchitis	204	25.8	10	1.3	63	12.6	10	2	53	7.6	6	1.3	320	16.1	26	1.5
Bronchiolitis/RSV	162	20.5	14	1.8	8	1.6	4	0.8	29	4.1	2	0.4	199	10	20	1.2
Any Listed	419	53	54	6.9	133	26.7	46	9.1	264	37.8	31	7	816	41	131	7.6

6 *Socioeconomic status was derived from a combination of mother's education level, health insurance status, and
7 household income weighted by region, see **Text A in S1 File** for more information.

8 **Table C.** Odds ratios and confidence intervals from the population-specific and combined association analysis between
9 respiratory illnesses in the first two years of life and physician-diagnosed asthma in the three main populations in GALA II
10 and SAGE: 2006-2014.

	Puerto Rican (n=1,574)	Mexican American (n=1,002)	African American (n=1,145)	Combined (n=3,721)
URI	10.6 (7.35-15.6)	3.71 (2.17-6.59)	10.2 (6.19-17.9)	8.45 (6.5-11.1)
Pneumonia	15.3 (6.77-44)	4.25 (2.23-8.82)	3.07 (1.73-5.82)	5.38 (3.66-8.2)
Bronchitis	29.4 (16.2-60)	6.56 (3.43-13.9)	5.42 (2.47-14.3)	14.7 (9.8-23.1)
Bronchiolitis/RSV	15.2 (8.98-27.8)	1.75 (0.531-6.77)	10 (2.96-62.7)	12.2 (7.56-21)
Any Listed	16.1 (11.8-22.2)	4.39 (2.98-6.6)	7.79 (5.31-11.8)	9.68 (7.84-12)

11 **BOLD** values are statistically significant at $p < 0.05$.

12 **Table D.** Pairwise comparison of odds ratios for asthma following early-life respiratory
 13 infection by racial/ethnic group including those diagnosed with asthma before age two.

Illness	Group	Group Pair		
		<i>African American</i>	<i>Combined</i>	<i>Mexican</i>
<i>URI</i>	<i>Combined</i>	0.537	-	-
	<i>Mexican</i>	0.010	0.009	-
	<i>Puerto Rican</i>	0.913	0.344	0.002
<i>Pneumonia</i>	<i>Combined</i>	0.129	-	-
	<i>Mexican</i>	0.483	0.561	-
	<i>Puerto Rican</i>	0.004	0.040	0.028
<i>Bronchitis</i>	<i>Combined</i>	0.042	-	-
	<i>Mexican</i>	0.735	0.051	-
	<i>Puerto Rican</i>	0.002	0.080	0.002
<i>Bronchiolitis/RSV</i>	<i>Combined</i>	0.802	-	-
	<i>Mexican</i>	0.073	0.004	-
	<i>Puerto Rican</i>	0.601	0.572	0.002
<i>Any Listed</i>	<i>Combined</i>	0.347	-	-
	<i>Mexican</i>	0.045	0.001	-
	<i>Puerto Rican</i>	0.005	0.009	0.000

15 **Table E.** Descriptive characteristics of Puerto Rican Islanders and Mainlanders.

	Islander				Mainlander			
	Case		Control		Case		Control	
	N	%	N	%	N	%	N	%
Number of subjects	237	100	725	100	47	100	58	100
Males	127	53.6	333	45.9	23	48.9	28	48.3
Underweight at birth	28	11.8	57	7.9	4	8.5	4	6.9
In-utero smoke exposure	7	3	33	4.6	7	14.9	7	12.1
Breastfed	132	55.7	410	56.6	9	19.1	22	37.9
Number of older siblings								
0	55	23.2	179	24.7	19	40.4	21	36.2
1	75	31.6	252	34.8	17	36.2	22	37.9
2 or more	107	45.1	294	40.6	11	23.4	15	25.9
Socioeconomic status*								
high	80	33.8	253	34.9	9	19.1	18	31
medium	46	19.4	128	17.7	5	10.6	8	13.8
low	111	46.8	344	47.4	33	70.2	32	55.2
Recruitment region								
Chicago	-	-	-	-	18	38.3	23	39.7
Houston	-	-	-	-	1	2.1	-	-
New York	-	-	-	-	27	57.4	34	58.6
San Francisco Bay Area	-	-	-	-	1	2.1	1	1.7
Puerto Rico	237	100	725	100	-	-	-	-
URI	50	21.1	24	3.3	7	14.9	11	19
Pneumonia	12	5.1	4	0.6	1	2.1	1	1.7
Bronchitis	35	14.8	9	1.2	1	2.1	1	1.7
Bronchiolitis/RSV	28	11.8	13	1.8	3	6.4	1	1.7

16 *Socioeconomic status was derived from a combination of mother's education level,
17 health insurance status, and household income weighted by region, see **Text A in S1**
18 **File** for more information.

Table F. Odds ratios and confidence intervals from the geography-specific analysis between respiratory illnesses in the first two years of life and physician-diagnosed asthma after the age of two in the two Puerto Rican populations: Islanders and Mainlanders.

	Islander (n=962)	Mainlander (n=102)
URI	7.8 (4.69-13.3)	0.817 (0.25-2.57)
Pneumonia	8.92 (3.04-32.4)	1.06 (0.0379-29.8)
Bronchitis	14.8 (7.23-33.4)	0.63 (0.0215-18.6)
Bronchiolitis/RSV	7.6 (3.9-15.6)	3.86 (0.386-91.3)
Any Listed	8.64 (5.72-13.2)	0.684 (0.222-2.01)

BOLD values are statistically significant at $p < 0.05$.