

Appendix A

Table A1. Hazard Ratios (HR) with 95% confidence interval (CI) for outcomes of Incidence fracture according to the GRS group, adjusted for FRAX score: results from sensitivity analysis after excluding subjects who received intervention in either of the three clinical trials. (N=14,722)

	Major osteoporotic fracture	Hip fracture
	HR (95 % CI)	HR (95 % CI)
Adjusted for FRAX probability		
low	1(reference)	1(reference)
medium	1.37 (1.11 - 1.70)	1.32 (0.98 - 1.77)
high	1.39 (1.10 - 1.74)	1.41 (1.03 -1.95)
Adjusted for FRAX probability + race		
low	1(reference)	1(reference)
medium	1.15 (0.93 - 1.43)	1.04 (0.77 - 1.40)
high	1.18 (0.93 -1.50)	1.15 (0.82 -1.61)

Significant results are in boldface.

Table A2. Hazard Ratios (HR) with 95% confidence interval (CI) for outcomes of Incidence fracture according to the GRS group, adjusted for FRAX score: results from sensitivity analysis after excluding subjects who received intervention in either of the three clinical trials. (N=14,722)

	Major osteoporotic fracture	Hip fracture
	HR (95 % CI)	HR (95 % CI)
Adjusted for FRAX probability		
Caucasian	1(reference)	1(reference)
American Indian	0.31 (0.19 -0.50)	0.32 (0.16 – 0.65)
Asian	0.10 (0.04 - 0.28)	0.15 (0.05 - 0.48)
AA	0.22 (0.18 - 0.27)	0.21 (0.16 - 0.28)
Hispanic	0.34 (0.27 - 0.43)	0.22 (0.15 - 0.31)
Adjusted for FRAX probability + GRS group		
Caucasian	1(reference)	1(reference)
American Indian	0.30 (0.19 - 0.49)	0.31 (0.15 – 0.64)
Asian	0.10 (0.04 - 0.27)	0.15 (0.05 - 0.47)
AA	0.23 (0.18 - 0.28)	0.22 (0.17 - 0.29)
Hispanic	0.34 (0.27 - 0.43)	0.21 (0.15 - 0.30)

Significant results are in boldface.

Figure A1. Box plot for the distribution of GRS in different race groups (N=23,981).

