

Indications that *Chymotrypsin-like Elastase 1* is Involved in Emphysema-**Supplement**

SUPPLEMENTAL METHODS

Animal Use

Animal Housing

Animal use was approved by the CCHMC Institutional Animal Use and Care Committee (2017-0064). Mice were housed in a pathogen-free facility with 12-hour light/dark cycles and provided chow and water *ad lib*. Previously generated *Cela1^{-/-}* (*1*) and wild type mice were all on the C57BL/6 background and derived from our existing colony.

Porcine Pancreatic Elastase Model of Emphysema

A single dose of 2 units porcine pancreatic elastase (PPE, Sigma, St. Louis, MO) at a concentration of 10 units/mL diluted in PBS was administered by tracheal instillation as previously described (2) to 8-12 week old C57BL/6 mice anesthetized with isoflurane and tracheally cannulated using an 18 gauge angiocatheter.

Aged Mouse Model of Emphysema

WT and *Cela1*^{-/-} mice were collected at age 70-75 weeks for evaluation of age-dependent alveolar simplification.

Mouse Lung Tissue Collection and Processing

At predetermined time points, mice were anesthetized with 0.2 mL of ketamine/xylazine/acepromazine and sacrificed by exsanguination. The left lung was ligated before inflation and used for protein and RNA analysis, and the right lung inflated at 25 cm H₂O water pressure with 4% PFA in PBS, fixed, paraffinized, lobes as previously described (?), and 5 μm sections created.

Mouse Lung Morphometry

Using the methods of Dunnill (3) on five images from each right lung lobe, mean linear intercepts were determined and used for comparisons.

anti-CELA1 Antibody Generation

100 micrograms of GEHNLSQNDGTEQYVNVQKIVSHPY (Genscript, Piscataway, NJ) peptide in 1 mL of PBS and 1 mL of Freund's complete adjuvant was administered subcutaneously at multiple sites to a New Zealand female rabbit with a subsequent administration of 100 and then 50 micrograms in incomplete Freund's adjuvant on days 21 and 42. Titers were determined by direct ELISA using CELA1-coated plates and titers of <1:5000 considered positive. 7 mL/kg of blood was collected every 2 weeks by marginal vein catheterization using isoflurane anesthesia.

Human Lung Tissue Use

Human tissue utilized under a waiver from the CCHMC IRB (2016-9641). Emphysematous lung explants from individuals with COPD and from aged individuals with no documented lung disease were obtained from the NHLBI lung tissue research consortium (LTRC). "Healthy" lung

specimens from non-lung organ donors were obtained from National Jewish Health Human Lung Tissue Consortium in Denver, Colorado. Flash frozen lung specimens of COPD, non-lung organ donors, and aged lung without known lung disease were obtained from the NIH lung tissue consortium and the National Jewish Health Human Lung Tissue Consortium. Portions of specimens were fixed and sectioned and other portions used for biochemical assays.

Biochemical Assays

Biochemical Assays

Left mouse lungs were ligated and collected prior to inflation and fixation of the right lungs. Protein and RNA was extracted from these specimens and used for Western Blot and PCR. Protein and RNA was similarly extracted from human lung specimens but additionally homogenized lung specimens were analyzed using Enzchek elastase, gelatinase, and proteinase assays (Thermo Fisher E12056, E12055, E6639).

Enzymatic Assays

Frozen human lung specimens were homogenized in RIPA buffer and protein content quantified. Ten μ g of protein was used for Enzcheck elastase, gelatinase, and proteinase assays per manufacturer instructions using a Molecular Devices Spectramax M2 plate reader using 4 hour readings for comparisons.

Proximity Ligation *in situ* Hybridization (PLISH)

Using oligos in Table 1 and previously published methods (4), PLISH for mouse *Cela1* human *CELA1* mRNA was performed. Briefly, sections were incubated with right and left sided oligonucleotides (Table 1), then linking oligonucleotides, ligation performed with T4 DNA ligase (New England Biolabs, M0202L). The sequences amplified by rolling circle amplification using Phi29 polymerase (New England Biolabs M0269L), and these oligos detected using a de-

tection oligonucleotide. DAPI counterstaining was performed and sections imaged on a Nikon NiE microscope.

Immunofluorescence

Human lung sections were incubated overnight with 1:500 dilution rabbit anti-CELA1 and 1:500 guinea pig anti-SCGB1A1 antibodies (gift of Jeffrey Whitsett) in 5% donkey serum with subsequent incubation with 1:5000 fluorophore-conjugated secondary antibodies, counterstained with DAPI, and mounted in prolong gold. Images were obtained using a Nikon NiE microscope.

Immunohistochemistry

Human lung sections were immunostained for CELA1 using anti-CELA1 guinea pig antibody (*I*) with secondary alone control using the ABC Vectastain kit (Vector Labs, Burlingame, CA). 4X tile scanned and 20X images were obtained using a Nikon 90i inverted microscope. Using Nikon elements software, the number of Cela1-positive cells per lung section were morphometrically determined in 4x tile scanned sections.

Western Blot

Mouse lung homogenates were electrophoretically separated, transferred to PVDF membranes, and total protein quantified using a total protein stain (LICOR, 926-11011). Blots were immunostained with anti-CELA1 guinea pig antibody (1:5,000 dilution) and anti-tropoelastin antibody (ab21600, Abcam, 1:5,000 dilution), anti-p16INK (Sigma SAB45000-72, 1:500), p19ARF antibody (Novus Biologics, NB200-169, 1:500), anti-p21 antibody (Novus Biologics, NBP2-29463, 1:500), anti-p53 antibody (Abcam, ab131442, 1:500) and evaluated by densitometry using an Odyssey system (LI-COR Biotechnology, Lincoln, NE). Fold-change values from total protein (ReVERT 700 Total Protein Stain, LI-COR Biotechnology) normalized values was

Table 1: PLISH Oligonucleotides

Oligo	Sequence
MsCela1_705_L243	TAGCGCTAACAACTTACGTCGTTATGAAGTCATGGGGCTGTCCACA
MsCela1_705_L742	TAGCGCTAACAACTTACGTCGTTATGTTCTGGCGACATTACAGCC
MsCela1_705_L130	TAGCGCTAACAACTTACGTCGTTATGTGAGACGGCCAA-GAGTTCCT
MsCela1_705_L552	TAGCGCTAACAACTTACGTCGTTATGTATAGGA-CACGCTGGGCAGG
MsCela1_705_L1009	TAGCGCTAACAACTTACGTCGTTATGTTTGCCCCAGATAC-CAAGAC
MsCela1_705_R243	CTCTCCGACAACCACTCGATTTATACGTC-GAGTTGAACGTCGTAACA
MsCela1_705_R742	CTCTGGTGAAGACGGTGGGCTTATACGTC-GAGTTGAACGTCGTAACA
MsCela1_705_R130	ACTGGTACTGGAGGGAAATCTTATACGTC-GAGTTGAACGTCGTAACA
MsCela1_705_R552	AGAGGAGCTGGAGCAGATGCTTATACGTC-GAGTTGAACGTCGTAACA
MsCela1_705_R1009	CTTTTCATTAACCCCCACCTTATACGTCGAGTTGAACGTCG-TAACA
HuCELA1_705_L707	TAGCGCTAACAACTTACGTCGTTATGtggacacaaagctggctact
HuCELA1_705_L51	TAGCGCTAACAACTTACGTCGTTATGtgggtgctgtgccataaag
HuCELA1_705_L228	TAGCGCTAACAACTTACGTCGTTATGttctggtaatccacgcagtg
HuCELA1_705_L539	TAGCGCTAACAACTTACGTCGTTATGagtccacagagggcaggtaa
HuCELA1_705_L458	TAGCGCTAACAACTTACGTCGTTATGagcagggactgtgttagcc
HuCELA1_705_R707	ctggacacaaagctggctacTTATACGTCGAGTTGAACGTCGTAACA
HuCELA1_705_R51	cattggtttccggaaggtccTTATACGTCGAGTTGAACGTCGTAACA
HuCELA1_705_R228	cagccaccacgcggaaagtcTTATACGTCGAGTTGAACGTCGTAACA
HuCELA1_705_R539	ggagctggagcagatggcgtTTATACGTCGAGTTGAACGTCGTAACA
HuCELA1_705_R458	cttgccccagcctgtgatgtTTATACGTCGAGTTGAACGTCGTAACA
MsNeg control 705 1	TAGCGCTAACAACTTACGTCGTTATGttataaccgtgcttcttca
MsNeg control 705 2	TAGCGCTAACAACTTACGTCGTTATGatccatgectgttttacaga
MsNeg control 705 3	cttcagatacgcctgattcTTATACGTCGAGTTGAACGTCGTAACA
MsNeg control 705 4	atgctttcatttacacagatTTATACGTCGAGTTGAACGTCGTAACA
Common Connector-Tye705	TGTTAGCGCTAACAAAATGCTGCTGCTGTACTACGAACAA-CAATACACATGTTACGACGT
Common Bridge	TCAACTCGACGTATAACATAACGACGTAAGT
Tye705 Probe	Tye705-ATGCTGCTGCTGTACTACG

used for calculations.

PCR

RNA was extracted from lung homogenates using RNEasy Mini columns (Qiagen, Valencia, CA) and cDNA library synthesized using a High Capacity cDNA Reverse Transcriptase Kit (Applied Biosystems, Foster City, CA). For human specimens, Taqman PCR was performed using the primers listed in Table 2 and a QuantStudio6 device (all Applied Biosystems). For mouse specimens Sybr Green PCR was performed using the primers in Table 3 and PowerUp SYBR Green (Applied Biosystems, AB25780).

Table 2: Taqman Primers

Target	Primer Catalog Number
Human Matrix Metaloproteinase-2	4331182_Hs01548727
Human Matrix Metaloproteinase-8	4453320-Hs01029057
Human Matrix Metaloproteinase-9	4453320-Hs00957562_m1
Human Matrix Metaloproteinase-12	4448892-Hs00159178
Human Matrix Metaloproteinase-14	4448892-Hs01037003
Human Proteinase-3	4331182-Mm00478323_m1
Human Cathepsin G	4331182-Mm00456011_m1
Human Neutrophil Elastase	4331182-Hs00236952_m1
Human Chymotrypsin-like Elastase 1	4331182- Hs00608115_m1
Eukaryotic 18S RNA	4333760T

Table 3: Sybr Green PCR Primers

Name	Sequence
MsCela1 FL1-Fwd	TTGTCCGAGAGCACAACCTG
MsCela1 FL1-Rev	CCAAGACACCAGCAGCATTC
MsGapdh(66-323) mRNA-F	AGAGTGTTTCCTCGTCCCGT
MsGapdh(66-323) mRNA-R	TGATGTAGTGGGGTCTCGC

Human *Ex vivo* Lung Stretch

Ten mm cores of frozen human lung tissue were created using a coring device and hand-cut 100-200 μ m sections were cut using a scalpel on dry ice. These sections were mounted onto a silicone mold and then subjected to biaxial stretch or imaged sequentially over time using a previously published 3D-printed confocal microscope-compatible stretching device (1,5,6) with Enzchek elastin zymography substrate (10 μ g/mL), Texas-Red conjugated albumin (Thermo Fisher A23017, 1 μ g/mL) and AF647-conjugated CELA1 (1 μ g/mL) (1). A 100 μ m Z-stack at 10x magnification was obtained using a Nikon A1 confocal microscope and the signal for each analyte at sequential levels of biaxial stretch were quantified and normalized to tissue autofluorescence. The rate of elastase activity, albumin binding, and CELA1 binding per fold change in area or time equivalent was measured per section and sections assayed in triplicate. The average rate of change per specimen was used for comparisons.

Human Genomic Studies

CELA1 gene variants were downloaded from the Broad Institute (7) , dbSNP (8) and Ensembl (9). Functional significance of variants was predicted using SIFT and PolyPhen-2 (10, 11).

Statistical Methods

Using R version 3.5.3 (12) , the following packages were used for statistical comparisons and graphics generation: ggplot2 (13) , gridExtra (14) , cowplot (15) , ggplotify (16) , corrplot (17) , and ggsignif (18). Welch's t-test and Welch's t-test and Wilcoxon rank-sum test were used to compare parametric and non-parametric data respectively. Parametric data is displayed as line and whisker plots with center line representing mean and whiskers standard deviation. Non-parametric data is displayed as violin plots with center diamond representing median value. For

both plot types dots represent individual data points. Correlative analyses were made by Pearson correlation coefficient. For all analyses p-values of less than 0.05 were considered significant.

SUPPLEMENTAL DATA

PLISH Negative Control

Figure S1. *Cela1* null and bacterial probe control images.




Figure 1: Negative Control Images for PLISH. (A) *Cela1*^{-/-} lung did not demonstrate any *Cela1*-mRNA at 21 days post-PPE. (B) A bacterial gene probe did not have any signal in wild type mouse lung at 21 days post-PPE.

Aged Mouse Lung Supplemental Data

There was no difference in *Cela1* protein levels in aged vs young WT mice (Figure S2A). Quantitative morphometry of Hart stained WT and *Cela1*^{-/-} lung specimens showed a trend towards more insoluble elastin in *Cela1*^{-/-} lungs (Figure S2A). There were no statistically significant differences in the amount of senescence-related proteins in aged WT and *Cela1*^{-/-} lung.




Figure 2: Additional Aged Mouse Lung Data. (A) Western blot of young (8-12 week-old) and aged (70-75 week-old) wild type mouse lung revealed no differences in Celsr1 protein levels. (B) Quantitative image analysis of elastin-stained mouse lung issues identified a trend towards increased total lung elastin in aged *Celsr1*^{-/-} mouse lungs. (C) Quantification of lung senescence proteins showed no differences except for a trend towards reduced p53 in aged *Celsr1*^{-/-} mouse lung.

CELA1 in Alveolar Type 2 Cells

We previously demonstrated that in AAT-deficient emphysema, CELA1 mRNA was present in a subset of alveolar type 2 (AT2) cells (19). In normal human lung and in COPD, we also found CELA1 protein in AT2 cells (Figure S3)

Aged Human Lung Supplemental Data

Western blot for CELA1 in aged smoker and non-smoker lung homogenates.




Figure 3: CELA1 in Human AT2 Cells. CELA1 (red) was present in a subset of AT2 cells in normal (shown here) and COPD human lung. Green signal is staining by HT2-280 which labels the apical membrane of human AT2 cells.




Figure 4: Aged Human Lung Western Blot. (A) CELA1 and (B) total protein stain of smoker and non-smoker aged human lung.

Lung Stretching Supplemental Data

Image of 3D-printed confocal microscope stretching device (Figure S4A&B). Quantification of biaxial strain applied by device (Figure S6C).

Supplemental Human Genomics Data

Mutations in the CELA1 human genome were assessed for functional significance. Although not predicted to be functional by SIFT, rs74336876 confers a mutation at the CELA1 propeptide cleavage site, The unannotated mutation at chr12:51733775 is the most common predicted functional mutation. Both had allele frequencies of <1%.




Figure 5: 3D-printed Confocal Microscope Lung Stretching Device. (A) Four motors are housed within a plastic case that fits within the microscope mounting plate. A silicone mold (circle) is secured by four clips attached to the motors by Tyvek strips. (B) Image of lung before stretching and after stretching (inset). (C) Quantification of the increase in mold aperture area with number of stretch (steps). The steps used for imaging are shown.

Table 4: Ten Most Common CELA1 Mutations and Functional Predictions

Position	RSID	Transcript Conse- quence	Protein Conse- quence	Type	Functional Prediction	Allele Frequency
51737562	rs17860300	175A>G	Met59Val	Missense	Benign	0.2376
51723598	rs60311818	628dupC	Leu210Profs	Ter24Frameshift	Benign	0.2295
51737607	rs17860299	130C>T	Arg44Trp	Missense	Benign	0.1982
51739648	rs17860287	30G>C	Gln10His	Missense	Benign	0.09864
51723499	rs17860364	728A>G	Gln243Arg	Missense	Benign	0.09134
51739640		38C>T	Pro13Leu	Missense	Benign	0.00909
51739625	rs74336876	53G>A	Arg18His	Missense	Benign	0.007377
51735144	rs17860313	47G>A	Arg116His	Missense	Benign	0.001723
51735071		419delC	Ala140Valfs	Ter47Missense	Benign	0.0005204
51733775		478G>A	Ala160Thr	Frameshift	Loss-of-Function	0.0002924

References

1. M. Ojha, R. Joshi, J. Lewis, Q. Fan, B. M. Varisco, Chymotrypsin-Like Elastase 1 Is Required for Late Lung Remodeling in Mouse Porcine Pancreatic Elastase Model of Emphysema.
2. B. Suki, E. Bartolak-Suki, P. R. M. Rocco, *Methods Mol Biol.* **1639** (2017).
3. M. S. Dunnill, *Thorax* **17**, 320 (1962).
4. M. Nagendran, D. P. Riordan, P. B. Harbury, T. J. Desai, *Elife* **7** (2018).
5. S. M. Young, *et al.*, *J Appl Physiol* (1985) **118**, 921 (2015).
6. R. Joshi, *et al.*, *Faseb j* **30**, 590 (2016).
7. 1000 Genomes | A Deep Catalog of Human Genetic Variation.
8. Home - dbGaP - NCBI.
9. ClinVar.
10. PolyPhen-2: prediction of functional effects of human nsSNPs.
11. SIFT - Predict effects of nonsynonymous / missense variants.
12. R. C. Team, *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria, 2013).
13. H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag New York, 2016).
14. B. Auguie, *gridExtra: Miscellaneous Functions for "Grid" Graphics* (2017).

15. C. O. Wilke, *cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'* (2019).
16. G. Yu, *ggplotify: Convert Plot to 'grob' or 'ggplot' Object* (2019).
17. T. Wei, V. Simko, *R package "corrplot": Visualization of a Correlation Matrix* (2017).
18. C. Ahlmann-Eltze, *ggsignif: Significance Brackets for 'ggplot2'* (2019).
19. M. Ojha, R. Joshi, J. Lewis, Q. Fan, B. M. Varisco, Chymotrypsin-Like Elastase 1 Is Required for Late Lung Remodeling in Mouse Porcine Pancreatic Elastase Model of Emphysema.