

### Supplementary Material 3. Computation of the Alternative Healthy Eating Index (AHEI)-2010

The AHEI-2010 score has been published previously [1].

The computation in the NutriNet-Santé cohort included 10 out of 11 original components as the intake of trans fatty acids was not available. Each component ranges from 0 to 10. Criteria for minimum and maximum scores are shown in the table below. Intermediate scores are attributed proportionally to the intake (see an example for the vegetable component in footnote 1)

Component	Criteria for minimum score (0)	Criteria for maximum score (10)
Vegetables <sup>1</sup> , servings/d	0	≥ 5
Fruit <sup>2</sup> , servings/d	0	≥ 4
Whole grains, g/d	0	Women: 75 Men: 90
Sugar-sweetened beverages and fruit juice <sup>3</sup> , servings/d	≥ 1	0
Nuts and legumes <sup>4</sup> , servings/d	0	≥ 1
Red or processed meat <sup>5</sup> , servings/d	≥ 1.5	0
Long-chain (n-3) PUFA (EPA + DHA), mg/d	0	250
PUFA, % of energy	≤ 2	≥ 10
Sodium <sup>6</sup> , mg/d	Highest decile	Lowest decile
Alcohol <sup>7</sup> , drinks/d	Women: ≥ 2.5 Men: ≥ 3.5	Women: 0.5-1.5 Men: 0.5-2.0
<b>Total</b>	<b>0</b>	<b>100</b>

<sup>1</sup> Potatoes and tubers are not included in the vegetable component; 1 serving = 118.3 g. Intermediate scores for vegetable intakes are calculated as follows: score = vegetable intake (g/d)\*10/(5\*118.3). An intake of 300 g will therefore translate to a score of 5.1 for the vegetable component

<sup>2</sup> Fruit juice are not included in the fruit component; 1 serving = 118.3 g.

<sup>3</sup> 100%-fruit juices are included in the sugar-sweetened beverages and fruit juice component; 1 serving is 226.8 g

<sup>4</sup> 1 serving = 28.35 g

<sup>5</sup> 1 serving of red meat is 113.4g and 1 serving of processed meat is 42.5g

<sup>6</sup> Sex-specific cut-offs were computed for deciles

<sup>7</sup> 1 drink of alcohol contains 10 g of ethanol

### REFERENCE

- 1 Chiuve SE, Fung TT, Rimm EB, *et al.* Alternative Dietary Indices Both Strongly Predict Risk of Chronic Disease. *The Journal of Nutrition* 2012;**142**:1009–18. doi:10.3945/jn.111.157222