

SUPPLEMENTAL MATERIAL

for

Phenotyping genetic differences in aldehyde dehydrogenase 2

after an alcohol challenge in humans

Joseph R. White, BS¹⁺, Freeborn Rwere, PhD¹⁺, Xiaocong Zeng, MD¹, Leslie McNeil, MPH¹,
Kevin Zhou, BS¹, Martin Angst, MD¹, Che-Hong Chen, PhD², Daria Mochly-Rosen, PhD², Eric
R. Gross, MD, PhD^{1*}

Supplemental Methods:

Exclusion criteria for study: Exclusion criteria were age less than 21 years or greater than 85 years, weight greater than 225 pounds, pregnancy or breastfeeding, no prior history of alcohol consumption, medication use or current health conditions that contraindicate alcohol consumption, shortness of breath in the last month, history of severe allergic reaction to alcohol, and absence of alcohol-induced flushing after one standard drink of alcohol.

ALDH2 Site-Directed Mutant Construction, Protein Expression and Purification: The colonies obtained from site directed mutagenesis were transformed in competent BL21 (DE3) cells and transformants grown in LB media overnight. The mutated plasmids were extracted and purified (QIAprep Mini-Kit, Qiagen, Chatsworth, CA). Following plasmid purification, the nucleotide sequences of the mutated ALDH2 was determined to confirm the presence of the desired mutations and absence of unwanted base changes (Sequetech, Mountain View, CA).

Protein Expression and Purification: Expression of human mitochondrial wild-type ALDH2 and its site directed mutants (R101G, R114W and E504K) was performed in freshly transformed competent BL21 (DE3) *Escherichia coli* and grown in 0.5L of NZCYM media containing 0.1 mM kanamycin and 1% (w/v) glucose. After induction with 0.5mM IPTG for 16 hours, the cells were harvested, lysed, and purified using a nickel affinity column for the histidine tagged protein (GE Healthcare Sciences, MA). A 10% SDS gel was run on the purified proteins to check their purity. The average purified yield for wild type, R114W, and E504K ALDH2 protein was 130-150 µg/mL.

Supplemental Tables:

Supplemental Table 1. Primers used to sequence an ALDH2 intron region (rs4646777) and ALDH2 exons. Twelve primer sets were used, with forward and reverse primers listed.

Supplemental Table 2. Formulas used to calculate breath metabolites for ethanol, acetaldehyde, and acetic acid using selective ion flow mass spectrometry. The reaction ion reaction ration and mass used for detection of ethanol, acetaldehyde, and acetic acid are listed.

Supplemental Table 3. Oligonucleotide primer sequences for generating ALDH2 point mutations from wild type ALDH2 enzyme for biochemical studies. Forward and reverse primers are listed that were used to sequence R101G, R114W and E504K.

Supplemental Table 4. Human volunteer demographics for study. Number of people, age, weight, sex and smoking status were reported for all volunteers recruited for the alcohol challenge portion of the study. No significant differences were noted between groups for any of these parameters.

Supplemental Figure Legends:

Supplemental Figure 1. Duration of tachycardia after alcohol challenge. **A.** Calculated acetaldehyde aldehyde area under the curve values for alcohol challenge. **B.** Duration of tachycardia during alcohol challenge. **C.** Change in heart rate from baseline versus peak acetaldehyde level 30 minutes after the start of the alcohol challenge. Blue data points are ALDH2*1*1 subjects and red data points are ALDH2*1*2 subjects. *=P<0.01 versus ALDH2*1*1 subjects, n=8/group (4 male and 4 female).

Supplemental Figure 2. Self-reported lineage of non-East Asian participants whose genotype was heterozygous for ALDH2*2. Participants who self-identified as non-East Asian were genotyped and 5 participants were identified to carry the ALDH2*2 genetic variant. These participants self-reported lineage from Mexico, Cuba, Iran, Europe, and Philippines. None reported ancestry from East Asian countries which have highly prevalent populations carrying the ALDH2*2 genetic mutation.

Supplemental Figure 3. Breath metabolite and physiologic characteristics after alcohol challenge in self-reported non-Asian subjects which are ALDH2*1*2. Breath concentrations (in parts per million) of **A.** ethanol **B.** acetaldehyde and **C.** acetic acid. Physiologic characteristics of subjects including **D.** heart rate **E.** skin temperature and **F.** mean arterial pressure. After collecting a baseline measurement, the next 2 measurements were during alcohol consumption. Subjects were monitored for an additional 140 minutes after alcohol consumption. Non-Asian subjects with the ALDH2*1*2 variant are represented with the black line. Blue lines are ALDH2*1*1 subjects and red lines ALDH2*1*2 subjects for comparison as previously reported in Figure 2. One subject was a heterozygote for both rs4646777 and ALDH2*1*2, *=P<0.01 versus ALDH2*1*1 subjects, n=4.

Supplemental Figure 4. Additional metabolic and physiologic data for R114W and R101G alcohol challenge testing. **A and B.** Ethanol levels measured by breath. **C and D.** Acetic acid levels measured by breath. **E and F.** Mean arterial pressure.

Supplemental Figure 5. Additional biochemical data for R114W relative to wild type ALDH2 and ALDH2 E504K using recombinant protein. **A.** Recombinant protein for wild type ALDH2, ALDH2 R114W, or ALDH2 E504K were used in an enzymatic activity assay that measures conversion of NAD⁺ to NADH based on relative absorbance. **B.** Enzymatic activity for purified protein from wild type ALDH2, ALDH2 R114W, and ALDH2 E504K. n=3 replicates/group, *P<0.01 relative to wild type ALDH2 enzyme activity.

Supplemental Table 1.

PCR Primer Sets for ALDH2 Exon Genotyping

Exon	Forward Primer	Reverse Primer
1	TAGCGCCACCCGCTTCGCTTGCATCA	TGCGGGGGACTCGGGCCGGAAAACAAA
2	GGTAGTCAGTATTAGGTTGACAGCTGG	TTAACACCGTTGTCTGCAGACAAG
3	TGTGCAGCGATATGCTGATGACC	GCTGCGAGCCTATTCACCACAT
4	TTGGAGAGACCATGGCAATAGTCCAG	TTACCTCCTAACAACGTTGCCTCCC
5	CAGAAAGACTCAGCTGGACCAGTTTG	TTGTCAGAGCCCATCTTCTTACCTGCC
6	CCAGTGTAGTTCTCTGAGGAAGCT	TAAGAGGGAGCCACTCTGGTTCACAT
7	GACCACATGTGTCCTTGGCAGA	GGAGCTCACCTTGAGCATGTCGT
8	TCTCTCGTGGTCCAGTTGCTCACT	ACAGCAACCAACGCCATTGGGCA
9	AGCAGGCATCAACCCTTACAGT	TTCTCATGCTGGCTCTAGACA
10	GGCTGCATAATTCTAAGCCTGAAGCCT	CATCAGGATGCCAGGCTGAACTGTTT
11	TTCCCCTGGAAGTGTTAGAGCATGGCT	ATTCCAGGATGGTGACCACCAGATTC
12	GTCCTGGGAGTGTAACCCATAA	AACAGACCCCAATCCCCCAGCA

Supplemental Table 2.

Metabolite	Reagent Ion	Reaction Ratio	Mass (m/z)
Ethanol	H ₃ O ⁺	2.7 x 10 ⁻⁹	47, 65
	NO ⁺	1.2 x 10 ⁻⁹	45, 63
Acetaldehyde	H ₃ O ⁺	3.7 x 10 ⁻⁹	45, 63
	NO ⁺	6.0 x 10 ⁻⁹	43, 61
Acetic Acid	H ₃ O ⁺	2.6 x 10 ⁻⁹	61, 79
	NO ⁺	9.0 x 10 ⁻¹⁰	90

Supplemental Table 3.


Mutant	Sequence
R101G (Forward)	5'- GAC GCA TCA CAC GGG GGC CGG CTG CTG -3'
R101G (Reverse)	5'- CAG CAG CCG GCC CCC GTG TGA TGC GTC -3'
R114W (Forward)	5'-CTG ATC GAG CGG GAC TGG ACC TAC CTG GCG-3'
R114W (Reverse)	5'-CGC CAG GTA GGT CCA GTC CCG CTC GAT CAG-3'
E504K(Forward)	5'-GGG CTG CAG GCA TAC ACT AAG GTG AAA ACT GTC-3'
E504K (Reverse)	5'-GAC AGT TTT CAC CTT AGT GTA TGC CTG CAG CCC-3'

Supplemental Table 4.


	*1/*1 East Asian	*1/*2 East Asian	Self-identified non-East Asian
Number of participants	8	8	10
Age (years)	28±2	29±2	33±4
Weight (pounds)	147±8	146±8	147±9
Sex (Male and Female)	4 and 4	4 and 4	2 and 8
Smoker	None	None	None

Supplemental Figure 1.


A.



B.




C.







Supplemental Figure 2.



Supplemental Figure 3.




Supplemental Fig 4.



Supplemental Fig 5.

A.



B.

