1 Supplementary Tables

2 Supplementary table 1

Group	Representative	Substituted	SNPs				
number	SNP						
1	rs6323	rs2235186	rs1137070	rs2064070	rs1465107	rs5906957	rs2072743
	T: 1194	G: 1194	C: 1187	T: 1176	G: 1178	G: 1286	C: 1074
	GT: 405	GA: 405	TC: 407	AT: 412	AG: 408	AG: 368	CT: 432
	G: 374	A: 374	T: 379	A: 385	A: 387	A: 319	T: 467
2	rs4923460	rs4923457	rs4074134	rs6484320	rs6265	rs1013442	rs1304100
	G: 1197	A: 1197	C: 1198	A: 1178	C: 1267	A: 1119	A: 1103
	GT: 686	TA: 686	CT: 685	TA: 695	CT: 637	AT: 732	GA: 745
	T: 90	T: 90	T: 90	T: 100	T: 69	T: 122	G:125
3	rs3794808	rs4583306	rs140701	rs1042173			
	C: 605	A: 625	C: 619	A: 550			
	TC: 975	GA: 970	CT: 973	CA: 983			
	T: 393	G: 378	T: 381	C: 440			
4	rs1360780	rs9470080	rs3800373	rs4713916			
	C: 941	C: 895	A: 1022	G: 1015			
	CT: 838	CT: 853	CA: 787	AG: 790			
	T: 194	T: 225	C: 164	A: 168			
5	rs2020936	rs6354	rs8076005				
	A: 1334	T: 1339	A: 1323				
	AG: 571	GT: 565	GA: 576				

	G: 68	G: 69	G: 74		
6	rs3769671	rs12914008	rs11030103		
	A: 1862	G: 1832	A: 1937		
	CA: 108	GA: 140	GA: 0		
	C: 3	A: 1	G: 36		
7	rs140700	rs11661134	rs4663866		
	C: 1671	G: 1720	A: 1675		
	TC: 292	GA: 244	CA: 285		
	T: 10	A: 9	C:13		
8	rs1051730	rs16969968			
	G: 798	G: 801			
	AG: 909	GA: 906			
	A: 266	A: 266			
9	rs4633	rs4680			
	C: 508	G: 508			
	CT: 992	GA: 993			
	T: 473	A: 472			
10	rs12702514	rs13222291			
	C: 638	G: 632			
	TC: 950	GA: 955			
	T: 385	A: 386			
11	rs10482605	rs6198			
	A: 1358	T: 1365			

	AG: 554	TC: 552			
	G [.] 61	C: 56			
	0.01	0.00			
10	*01040710	ro10654779			
12	rs1042713	1512654778			
	0.700	0.740			
	G: 730	G: 749			
	AG: 937	GA: 928			
	A: 306	A: 296			
13	rs4818	rs6269			
	C: 705	A: 676			
	GC: 933	GA: 946			
	C: 335	G: 351			
	9. 333	6.331			
44					
14	151360/80	189470080			
	_	-			
	T: 1328	C: 1359			
	TA: 587	CT: 558			
	A: 58	CT: 56			
15	rs10521432	rs6651806			
	G: 1197	A: 1144			
	GA: 384	CA: 405			
	Δ· 302	C: 424			
	N. 002	0. 424			
16	re10925210	rc11020101			
10	1510035210	1511030101			
	0.000				
	C: 620	A: 531			
	CA: 951	TA: 992			
	A: 402	T: 450			

Supplementary table 1: Groups of SNPs used in this study. Groups of correlated SNPs as a result of linkage disequilibrium analysis. SNPs were assigned to the same group if a correlation of $r^2 \ge 0.5$ with the group's representative was calculated. The representative SNP for each group (with the highest r^2 -values for intergroup correlations) is indicated in bold. Abbreviations: T: thymine, C: cytosine, G: guanine, A: adenine, SNP: single nucleotide polymorphism.

11 Supplementary table 2

Models for SNP screening

Control variables	Univariable	Multivariable without variable	Multivariable with variable
		elimination	elimination
			ennindien
Sex: male	No	Yes	No
Diagnosis	No	Yes	No
Time since diagnosis (vears)		Yes	Yes
····· () ···· ()			
Age (years)	No	Yes	Yes
		X	
ВМІ	No	Yes	Yes
Smoking status	No	Yes	Yes
Daily alcohol consumption	No	Yes	Yes
Disease activity (standardized	No	Yes	Yes
CDAI/MTWAI)			
Disease related surgery prior to	No	Yes	No
enrollment			
Systemic steroids	No	Yes	No
INF INHIBITORS	No	Yes	No
Number current therapies	No	Yes	Yes

Supplementary table 2: Different SNP screening models. This table shows which control variables were part of the models used to screen for depression-associated SNPs. Abbreviations: BMI: body mass index, CDAI: Crohn's Disease Activity Index, MTWAI: Modified Truelove and Witts Severity Index, TNF: tumor necrosis factor.

17 Supplementary table 3

18

Cox proportional hazards model - hazard ratios for depression in IBD patients over time

	Endpoint variable: depression (HADS depression sub score ≥11)						
		Univariable	Multivariable				
rs588765: TC (vs. CC)	0.67 (CI: 0.45-1.02)		0.66 (CI: 0.44-1.01)				
	*		*				
rs588765: TT (vs. CC)	0.99 (CI: 0.60-1.63)		1.01 (CI: 0.61-1.68)				
rs2522833: AC (vs.		1.39 (CI: 0.91-		1.42 (CI: 0.93-			
AA)		2.11)		2.17)			
rs2522833: CC (vs.		1.19 (CI: 0.69-		1.13 (CI: 0.66-			
AA)		2.047		1.95)			
-							

19

Supplementary table 3: Predictors for depression (HADS-D ≥11) over time. Cox 20 proportional hazards models showing the predictive value of different SNP variables 21 for the occurrence of depression in IBD patients. Results are presented as hazard 22 23 ratio, control variables of the multivariable models are omitted; p-values are indicated 24 as follows: *: p<0.1, **: p<0.05, ***: p<0.01. 919 patients were analyzed and 114 times the endpoint was reached. Abbreviations: CI: 95% confidence interval, IBD: 25 inflammatory bowel disease, HADS: Hospital Anxiety and Depression Scale, T: 26 27 thymine, C: cytosine.

29 Supplementary table 4

Cox proportional hazards model - hazard ratios for clinical deterioration in IBD patients over time

	Endpoint variable: active disease (CDAI/MTWAI)						
		Univariable	Multivariable				
rs588765: TC (vs. CC)	0.74 (Cl: 0.53-		0.74 (CI: 0.52-1.06)				
	1.07)		*				
rs588765: TT (vs. CC)	1.13 (Cl: 0.74-		1.10 (CI: 0.72-1.69)				
	1.72)						
rs2522833: AC (vs.		1.05 (Cl: 0.73-		1.05 (CI: 0.73-			
ΑΑ)		1.50)		1.50			
rs2522833: CC (vs.		1.28 (Cl: 0.83-		1.28 (Cl: 0.83-			
AA)		1.96)		1.96)			

30

31 Supplementary table 4: Predictors for active disease in IBD patients over time.

32 Cox proportional hazards models showing the predictive value of different variables for the occurrence of active disease measured by CDAI and MTWAI in IBD patients. 33 34 Results are presented as hazard ratio, control variables of the multivariable models are omitted; p-values are indicated as follows: *: p<0.1, **: p<0.05, ***: p<0.01. 1006 35 observations with 157 events were analyzed. Abbreviations: CI: 95% confidence 36 interval, IBD: inflammatory bowel disease, CDAI: Crohn's Disease Activity Index, 37 MTWAI: Modified Truelove and Witts Severity Index, A: adenine T: thymine, C: 38 cytosine. 39

41 Supplementary figure 1



42

43

Supplementary figure 1: Pairwise linkage disequilibrium matrix of SNPs. Graphic
 illustrating the correlation between each possible pair of SNPs measured by the
 square of the correlation coefficient (r²).



49

50

Supplementary figure 2: SNPs' association significances. Array of the p-values for each SNPs' five possible inheritance patterns association with depression at enrollment in an univariable model. The upper red dotted line indicates a Bonferronicorrected significance level of p < 0.05.



57

58

59 **Supplementary figure 3: SNPs' association significances.** Array of the p-values 60 for each SNPs' five possible inheritance patterns association with depression at 61 enrollment in the complete multivariable model. The upper red dotted line indicates a 62 Bonferroni-corrected significance level of p < 0.05.



64

Supplementary figure 4: SNPs' association significances. Array of the p-values for each SNPs' five possible inheritance patterns association with depression at enrollment in the reduced multivariable model. The upper red dotted line indicates a Bonferroni-corrected significance level of p < 0.05.



Supplementary figure 5: Rs2522833 genotype does not significantly impact on
hazard for depression and IBD activity over time. A: Percentage of IBD patients
without depression stratified for rs2522833 genotype (log-additive). B: Percentage of
IBD patients without active disease defined as CDAI ≥150 or MTWAI ≥10 in CD and
UC patients, respectively and stratified for rs2522833 genotype (log-additive).
Analysis: multivariable Cox proportional hazards models.



82 Supplementary figure 6: Depression increases the hazard for adverse outcomes

83 **over time.** Percentage of IBD patients without experiencing the respective outcome

- 84 (A-H) stratified for depression. Analysis: univariable Cox proportional hazards models.
- 85



Supplementary figure 7: Depression increases the hazard for EIMs over time.
Percentage of IBD patients without experiencing the respective outcome (A-H)
stratified for depression. Analysis: univariable Cox proportional hazards models.

92





97 Supplementary figure 8: Visualization of rs588765's and rs2522833's genotype 98 distribution regarding depression and active disease. A: Percentage of IBD patients without depression stratified for rs588765 genotype (codominant with 99 rs588765-CC as reference). B: Percentage of IBD patients without depression 100 101 stratified for rs2522833 genotype (codominant with rs2522833-AA as reference). C: 102 Percentage of IBD patients without active disease defined as CDAI ≥150 or MTWAI 103 ≥10 in CD and UC patients, respectively and stratified for rs588765 genotype (codominant with rs588765-CC as reference). D: Percentage of IBD patients without 104 105 active disease defined as CDAI ≥150 or MTWAI ≥10 in CD and UC patients, 106 respectively and stratified for rs2522833 genotype (codominant with rs2522833-AA as 107 reference).