

Additional File S4 - Mode and dynamics of *vanA*-type vancomycin-resistance dissemination in Dutch hospitals

Authors: Sergio Arredondo-Alonso¹, Janetta Top¹, Jukka Corander^{2,3,4}, Rob J L Willems¹, Anita C Schürch¹

¹Department of Medical Microbiology, University Medical Center Utrecht, Utrecht, The Netherlands.

²Pathogen Genomics, Wellcome Trust Sanger Institute, Cambridge CB10 1SA, UK.

³Department of Biostatistics, University of Oslo, Oslo, Norway.

⁴Department of Mathematics and Statistics, Helsinki Institute of Information Technology (HIIT), FI-00014 University of Helsinki, Finland.




Figure S1 Distribution of pairwise Mash distances ($k = 21$, $s = 1,000$) between *vanA* complete plasmid sequences ($n = 26$).




Figure S2. Heatmap and single-linkage clustering of the *pyani* pairwise alignment identity in the set of *vanA* complete plasmid sequences ($n = 26$).




Figure S3. Visualization of the multiple genome alignment of complete plasmid sequences (n = 7) belonging to the plasmid type B. A minimum blast length of 500 bp and minimum identity of 80% were considered to plot the alignment blocks.




Figure S4. Scheme of the gene structure and content present in the plasmid sequences to infer vancomycin-resistant plasmid types (A-F). For visualization purposes, only the genes belonging to the *vanA* cluster and IS elements present in the plasmids are indicated.




Figure S5. Distribution of pairwise Mash distances ($k = 21$, $s = 1,000$) between predicted *vanA* plasmid sequences ($n = 282$).




Figure S6. Contribution of nested genetic elements in the dissemination of vancomycin resistance in specific Dutch hospitals. VRE samples isolated in 12 consecutive months and from the same hospital were considered. Clonal dissemination (green bar) corresponded to pairs of isolates sharing the same hierBAPS SC, *vanA* plasmid type and Tn1546 variant. Plasmid

dissemination (orange bar) corresponded to pairs of isolates sharing the same *vanA* plasmid type and Tn1546 variant but different hierBAPS SC. Dissemination occurring by Tn1546 transposition (blue bar) corresponded to pairs of isolates sharing the same Tn1546 variant, different *vanA* plasmid type and same or different the hierBAPS SC. Unrelated (purple bar) isolates corresponded to pairs of isolates with a different Tn1546 variant.




Figure S7. Contribution of nested genetic elements in the dissemination of vancomycin resistance divided by years (2012-2013, 2013-2014 and 2014-2015). VRE samples isolated in 12 consecutive months and from the same hospital were considered. Clonal dissemination (green bar) corresponded to pairs of isolates sharing the same hierBAPS SC, *vanA* plasmid type

and Tn1546 variant. Plasmid dissemination (orange bar) corresponded to pairs of isolates sharing the same *vanA* plasmid type and Tn1546 variant but different hierBAPS SC. Dissemination occurring by Tn1546 transposition (blue bar) corresponded to pairs of isolates sharing the same Tn1546 variant, different *vanA* plasmid type and same or different the hierBAPS SC. Unrelated (purple bar) isolates corresponded to pairs of isolates with a different Tn1546 variant.