



SUPPLEMENTARY INFORMATION

ACE2 expression in adipose tissue is associated with COVID-19 cardio-metabolic risk factors and cell type composition

Julia S. El-Sayed Moustafa^{1*}, Anne U. Jackson^{2*}, Sarah M. Brotman^{3*}, Li Guan^{4*}, Sergio Villicaña¹, Amy L. Roberts¹, Antonino Zito^{1,5,6}, Lori Bonnycastle⁷, Michael R. Erdos⁷, Narisu Narisu⁷, Heather M. Stringham², Ryan Welch², Tingfen Yan⁷, Timo Lakka^{8,9,10}, Stephen Parker⁴, Jaakko Tuomilehto^{11,12,13}, Francis S. Collins⁷, Päivi Pajukanta¹⁴, Michael Boehnke², Heikki A. Koistinen^{15,16,17}, Markku Laakso^{18,19}, Mario Falchi¹, Jordana T. Bell¹, Laura J. Scott^{2**}, Karen L. Mohlke^{3**} and Kerrin S. Small^{1**}



Supplementary Figure 2: Age and BMI distributions in the TwinsUK, METSIM and FUSION studies. a-c) Age distribution in TwinsUK, METSIM and FUSION studies. d-f) BMI distribution in TwinsUK, METSIM and FUSION studies. Plots include only non-diabetic subjects.




Supplementary Figure 3: ACE2 expression levels across studies and tissues.

ACE2 expression is plotted as TMM-adjusted counts per million. a) TwinsUK adipose tissue b) METSIM study adipose tissue c) FUSION study adipose tissue d) TwinsUK skin.


Expression of *ACE* and *TMPRSS2* across TwinsUK tissues

We also assessed expression of *ACE2* in skin, lymphoblastoid cell lines (LCLs) and whole blood in TwinsUK. In matched samples from the same subjects in TwinsUK, *ACE2* expression was lower in skin than in adipose tissue (Supplementary Figure 3), and undetectable in lymphoblastoid cell lines and whole blood.


Beyond *ACE2*, two other components of the RAS system of interest are angiotensin-converting enzyme (*ACE*), and transmembrane protease, serine 2 (*TMPRSS2*). *ACE* plays a central role in regulation of blood pressure²⁵, while *TMPRSS2* has received attention since its identification as a cofactor aiding viral invasion by SARS-CoV-2⁷. We therefore explored their expression and correlation patterns in the TwinsUK multi-tissue sample. *ACE* expression was detected in all four tissues, while *TMPRSS2* expression passed our filters only in skin. When comparing normalised, non-transformed gene counts per million (TMM-adjusted CPMs; TMMAdjCPMs), *ACE* had higher expression levels than *ACE2*, with expression much lower in LCLs and whole blood compared to adipose tissue and skin (median[SD] *ACE* adipose = 41.65[14.02]; median[SD] *ACE* skin = 33.30[16.50]; median[SD] *ACE* LCL = 0.17[0.23]; median[SD] *ACE* whole blood = 3.53[1.70]) (Supplementary Figure 4). *TMPRSS2* expression in skin was also higher than that of *ACE2* (median [SD] *TMPRSS2* skin = 18.63[16.42]). We observed an inverse correlation between *ACE2* and *ACE* expression in adipose tissue (Spearman Rho = -0.19; $P = 8.40 \times 10^{-5}$) but not skin (Spearman Rho = -0.03; $P = 0.53$). No correlation was observed between *ACE2* expression in adipose tissue and skin in the same subjects (Spearman Rho = -0.06; $P = 0.23$). On the other hand, *ACE* expression in adipose tissue and skin was positively correlated (Spearman Rho = 0.27; $P = 2.21 \times 10^{-7}$), with no significant correlation across the remaining tissues (Supplementary Figure 4).




Supplementary Figure 4: Expression levels of RAS genes *ACE* and *TMPRSS2* across tissues in TwinsUK. a-e) Histograms of expression levels of *ACE* and *TMPRSS2* expression in TwinsUK skin, LCL whole blood, and adipose tissue, as indicated. f) Correlation plot of RAAS gene expression across tissues in unrelated subjects from the TwinsUK study (n=463 participants).




Supplementary Figure 5: *ACE2* expression in adipose tissue and skin show inverse direction of association with age. a) TwinsUK adipose tissue. b) TwinsUK skin. Each point represents an individual. Participant age was rank inverse-normal transformed. *ACE2* expression residuals plotted were adjusted for technical residuals and BMI to correspond to the reported association model.



Supplementary Figure 6: Correlation plot of adipose tissue *ACE2* expression, phenotypic variables, and adipose tissue estimated cell type compositions in TwinsUK. Adipose tissue *ACE2* expression residuals were adjusted for RNASeq technical covariates. Correlations were assessed in unrelated subjects (n=441).



Supplementary Figure 7: Adipose tissue DNA methylation at methylation probes proximal to the *ACE2* gene is not associated with *ACE2* expression levels. Each point represents a single subject, with adipose tissue methylation beta values plotted on the x-axis and adipose tissue *ACE2* expression on the y-axis. Rank-based inverse normal transformation was applied to both methylation and gene expression data.



Supplementary Figure 8: Estimated cell type proportion of whole blood in the METSIM and FUSION studies. Cell type proportions were estimated using CIBERSORT (METSIM) and DESeq2 (FUSION) (See methods).