

SUPPLEMENTARY INFORMATION

SUPPLEMENTARY NOTE:

- a. Members of the Japanese Research Committee on Idiopathic Osteonecrosis of the Femoral Head ----- P2
- b. Acknowledgement ----- P3
- c. Web resources ----- P5

SUPPLEMENTARY FIGURES:

Supplementary Figure 1: Quantile-quantile (QQ) plot of the meta-analysis association *P* values for all variants (left) and the variants outside of known SLE loci (right). ----- P6

Supplementary Figure 2: Regional association plots for the 113 SLE genomic loci. ----- P7

Supplementary Figure 3: Expression quantitative trait locus (eQTL) effect of rs66977652 on *SMG7* gene across a range of human tissues/cells in GTEx v8. ----- P28

Supplementary Figure 4: Tissues (a) and cells (b) in which the expression of genes within SLE-associated loci are significantly enriched in DEPICT analysis. ----- P29

Supplementary Figure 5: Schematic diagram of luciferase expression vector constructs. ----- P30

a. Member of the Japanese Research Committee on Idiopathic Osteonecrosis of the Femoral Head

Kohei Tomizuka¹, Taisuke Seki², Yoshiya Tanaka³, Toshikazu Kubo⁴, Ryo Hisada⁵, Tomokazu Yoshioka⁶, Mihoko Yamazaki⁷, Tamon Kabata⁸, Yoshiharu Amasaki⁹, Yoichi Ohta¹⁰, Takahiro Okawa¹¹, Yohei Naito¹², Ayumi Kaneuji¹³, Yuji Yasunaga¹⁴, Kenji Ohzono¹⁵

¹Laboratory for Statistical and Translational Genetics Analysis, RIKEN Center for Integrative Medical Sciences

²Department of Orthopaedic Surgery, Nagoya University Graduate School of Medicine

³The First Department of Internal Medicine, University of Occupational and Environmental Health, School of Medicine

⁴Graduate School of Medical Science Kyoto Prefectural University of Medicine

⁵Obihiro Kosei Hospital

⁶Division of Regenerative Medicine for Musculoskeletal System, Faculty of Medicine, University of Tsukuba

⁷Kido Hospital

⁸Department of Orthopaedic Surgery, Graduate School of Medical Science, Kanazawa University

⁹Kuriyama Red Cross Hospital, Department of Rheumatology

¹⁰Department of Orthopaedic Surgery, Osaka City University Graduate School of Medicine

¹¹Orthopedis and Joint Surgery Center, Kurume Univ. Medical Center

¹²Department of Orthopaedic Surgery, Mie University Graduate School of Medicine

¹³Department of Orthopaedic Surgery, Kanazawa Medical University

¹⁴Hiroshima Prefectural Rehabilitation Center

¹⁵Department of Orthopaedic Surgery, Amagasaki Chuo Hospital

b. Acknowledgement

We appreciate the contribution of following colleagues in the Japanese Research Committee on Idiopathic Osteonecrosis of the Femoral Head.

Yasuharu Nakashima¹, Hiroshi Kataoka², Makoto Kondo², Tsuyoshi Asano³, Tohru Irie³, Norimasa Iwasaki³, Tatsuya Atsumi⁴, Shunji Kishida⁵, Shigeo Hagiwara⁶, Yukihide Iwamoto⁷, Satoshi Ikemura⁷, Ryosuke Yamaguchi⁷, Kazuyuki Karasuyama⁷, Kazuhiko Sonoda⁷, Yusuke Kubo⁷, Takeshi Utsunomiya⁷, Hiroyuki Hatanaka⁷, Shuji Baba⁷, Koichiro Kawano⁷, Noriko Yamamoto⁷, Ichiei Narita⁸, Koichi Akashi⁹, Hiroshi Tsukamoto¹⁰, Yojiro Arinobu¹¹, Mitsuteru Akahoshi¹², Hiroki Mitoma¹¹, Masahiro Ayano¹³, Atsuhiko Fujie¹⁴, Yoshiaki Toyama¹⁴, Atsushi Funayama¹⁴, Kunihiro Yamaoka¹⁵, Hironari Hanaoka¹⁶, Katsunori Ikari¹⁷, Hisashi Yamanaka¹⁷, Tetsuji Hosozawa¹⁷, Shigeki Momohara¹⁷, Fumio Sekiya¹⁸, Masakazu Matsushita¹⁸, Megumi Matsuhashi¹⁹, Kazuhide Tanimura¹⁹, Takashi Sakai²⁰, Wataru Ando²¹, Takashi Nishii²¹, Masaki Takao²¹, Ryosuke Nakanishi²², Yukiharu Hasegawa²³, Kazuyoshi Saito²⁴, Kazuhisa Nakano²⁵, Tetsuya Sakamoto²⁶, Keiichiro Ueshima²⁷, Yoshitomo Kajino²⁸, Tomomichi Kajino²⁹, Hiroaki Nakamura³⁰, Shigekazu Mizokawa³⁰, Shinichiro Kume³¹, Akihiro Sudo³², Masahiro Hasegawa³², Toru Ichiseki³³, Takuma Yamasaki³⁴

¹Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kyushu University

²Department of Rheumatology & Clinical Immunology, Sapporo City General Hospital

³Department of Orthopaedic Surgery, Faculty of Medicine and Graduate School of Medicine, Hokkaido University

⁴Department of Rheumatology, Endocrinology and Nephrology, Faculty of Medicine and Graduate School of Medicine, Hokkaido University

⁵Department of Orthopaedic Surgery, Seirei Sakura Citizen Hospital

⁶Department of Orthopaedic Surgery, Graduate School of Medicine, Chiba University

⁷Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kyushu University

⁸Division of Clinical Nephrology and Rheumatology, Kidney Research Center, Niigata University Graduate School of Medical and Dental Sciences.

⁹Faculty of Medical Sciences, Kyushu University

¹⁰Shin-Kokura Hospital

¹¹Kyushu University Hospital

¹²Faculty of Medicine, Saga University

¹³Graduate School of Medical Sciences, Kyushu University

¹⁴Department of Orthopaedic Surgery, Keio University School of Medicine

¹⁵Department of Rheumatology and Infectious Diseases, Kitasato University School of Medicine

¹⁶Division of Rheumatology, Department of Internal Medicine, Keio University School of Medicine

¹⁷Institute of Rheumatology, Tokyo Women's Medical University, Tokyo, Japan

- ¹⁸Department of Internal Medicine and Rheumatology, Juntendo University School of Medicine
- ¹⁹Hokkaido Medical Center for Rheumatic Diseases
- ²⁰Department of Orthopaedic Surgery, Yamaguchi University Graduate School of Medicine
- ²¹Department of Orthopaedic Medical Engineering, Osaka University Graduate School of Medicine
- ²²Department of Orthopaedic Surgery, Fujigaoka Hospital, Showa University School of Medicine
- ²³Department of Rehabilitation, Kansai University of Welfare Science, Osaka, Japan
- ²⁴Tobata General Hospital
- ²⁵The First Department of Internal Medicine, University of Occupational and Environmental Health, Japan
- ²⁶Department of Orthopaedic Surgery, Faculty of Medicine, Fukuoka
- ²⁷Kyoto interdisciplinary institute hospital of community medicine
- ²⁸Department of Orthopaedic Surgery, Graduate School of Medical Science, Kanazawa University
- ²⁹Tonan hospital, Department of orthopaedic surgery
- ³⁰Department of Orthopaedic Surgery, Osaka City University Graduate School of Medicine
- ³¹Orthopedic and Joint Surgery Center, Kurume Univ. Medical Center
- ³²Department of Orthopaedic Surgery, Mie University Graduate School
- ³³Department of Orthopaedic Surgery, Kanazawa Medical University
- ³⁴Department of Orthopaedic Surgery, National Hospital Organization Kure Medical Center and Chugoku Cancer Center

c. Web resources

The software used has been described in details in Online Methods section. Software and resources included:

Pre-phasing: SHAPEITv2/v2.3, https://mathgen.stats.ox.ac.uk/genetics_software/shapeit

Pre-phasing: Eagle v2, <https://data.broadinstitute.org/alkesgroup/Eagle/>

Imputation: minimac3/4, Michigan Imputation Server,
<https://imputationserver.sph.umich.edu>

Imputation: IMPUTEv2/v4, <https://jmarchini.org/impute-4/>

Association: EPACTSv3.3 (EMMAX), <https://github.com/statgen/EPACTS>

Association: Plinkv1.9, <https://www.cog-genomics.org/plink/>

Association: SNPTESTv2.5, https://mathgen.stats.ox.ac.uk/genetics_software/snpTEST

Meta-analysis: metal, <https://genome.sph.umich.edu/wiki/METAL>

Conditional analysis: GCTAv1.9, <https://cnsgenomics.com/software/gcta/#Overview>

Bayesian statistical fine-mapping, FINEMAP v1.4, <http://www.christianbenner.com>

Visualization: LocusZoom, https://genome.sph.umich.edu/wiki/LocusZoom_Standalone

Visualization: UCSC genome browser, <https://genome.ucsc.edu>

Gene-based association: MAGMA, <https://ctg.cncr.nl/software/magma>

Gene-based association: FUMA, <https://fuma.ctglab.nl>

Gene-based association: FUSION, <http://gusevlab.org/projects/fusion/>

Gene-based association: DEPICT, <https://data.broadinstitute.org/mpg/depict/>

Heritability estimation: LDSC, <https://github.com/bulik/ldsc>

Cells/Tissues enrichment and Gene-set analysis: DEPICT,
<https://data.broadinstitute.org/mpg/depict/documentation.html>

Drug targets enrichment: GREP, <https://github.com/saorisakaue/GREP>

Mutation effect prediction: MENTR, <https://github.com/koido>

Regulatory element intersection: RELI, <https://github.com/WeirauchLab/RELI>

Resource: GTEx Portal, <https://gtexportal.org/home/>

ENCODE blacklist regions: [https://github.com/Boyle-](https://github.com/Boyle-Lab/Blacklist/blob/master/lists/hg19-blacklist.v2.bed.gz)

[Lab/Blacklist/blob/master/lists/hg19-blacklist.v2.bed.gz](https://github.com/Boyle-Lab/Blacklist/blob/master/lists/hg19-blacklist.v2.bed.gz)

Gene Expression Omnibus: <https://www.ncbi.nlm.nih.gov/geo/>

fastQC: <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>

samtools: <http://www.htslib.org>

bedtools: <https://bedtools.readthedocs.io/en/latest/>


bowtie2: <http://bowtie-bio.sourceforge.net/bowtie2/index.shtml>

PICARD: <https://broadinstitute.github.io/picard/>


MACS2: <https://github.com/macs3-project/MACS>

Trim Galore: https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/


Supplementary Figure 1. Quantile-quantile (QQ) plots of the meta-analysis association P values for all variants (left) and the variants outside of known SLE loci (right). X-axis is the minus $\log_{10}$ -transformed expected P values while the y-axis is the minus $\log_{10}$ -transformed observed P values in the meta-analysis. The genomic control inflation factor (λ_{gc}) is shown in the top of each figure.




region 11: rs13385731




region 12: rs11126034




region 13: rs10207954




region 14: rs73954925




region 15: rs218174




region 16: rs11679244




region 18: rs7572733




region 20: rs438613




region 30: rs4643809




region 32: rs7725218




region 33: rs2544920




region 34: rs370449198




region 35: rs2549002




region 37: rs10036748




region 38: rs2421184




region 39: rs2431697




region 40: rs9503037




region 41: rs9268807




region 45: rs548234




region 46: rs9488914




region 48: rs9322454




region 50: rs117026326




region 52: rs2736332




region 53: rs2272736




region 54: rs142937720




region 55: rs17374162




region 56: rs16902895




region 57: rs1887428




region 58: rs7858766




region 59: rs77448389




region 61: rs7902146




region 62: rs10995261




region 96: rs35966917




region 97: rs2671655




region 98: rs113417153




region 99: rs1788097




region 100: rs118075465




region 101: rs2238577




region 102: rs5826945




region 103: rs55882956




region 2: rs116785379 (1:157108159:G:C), rs112806509(1:157502015:AT:A), and rs2317230(1:157674997:G:T)




region 3: rs12120358(1:161444369:A:T), rs111994823(1:161470274:T:C), rs76107698(1:161569829:G:C), and rs75773410(1:161625798:A:G)




region 4: rs2205960 (1:173191475:G:T), rs117278480 (1:173210789:A:G), rs844646 (1:173216373:A:G), and rs2039982 (1:173359187:C:T)




region 5: rs13306575 (1:183532437:G:A) and rs66977652 (1:183540215:TA:T)




region 17: rs11889341 (2:191943742:C:T) and rs71030321 (2:191948986:T:TA)




region 19: rs7565158 (2:213593970:G:T) and rs2371790 (2:213884666:A:G)




region 27: rs6533951 (4:79644279:A:G) and rs2867700 (4:81084349:A:G)




region 31: rs956237 (4:109046960:G:A) and rs57634688 (4:109111805:C:T)




region 36: rs115267018 (5:133421109:G:C), rs6874758 (5:133429192:G:C), and rs138305363 (5:133714873:A:G)




region 42: rs111658701 (6:34228565:T:C), rs6457796 (6:34828553:T:C), and rs10484578 (6:35246319:A:G)




region 43: rs56285849 (6:36704293:T:C) and rs34868004 (6:36715031:C:CA)




region 44: rs3857496 (6:90719164:T:C) and rs597325 (6:91002494:A:G)




region 47: rs75163761 (6:138157489:C:T) and rs9494894 (6:138228520:T:C)




region 49: rs62447171 (7:50309890:G:A) and rs10239000 (7:50332196:A:G)




region 51: rs3757387 (7:128576086:T:C), rs41298401 (7:128578307:C:G), rs78724056 (7:128654051:G:T), and rs28364822 (7:128695982:T:A)




region 60: rs7072606 (10:49933974:T:C) and rs7097397 (10:50025396:G:A)




region 64: rs11599734 (10:103963397:C:G) and rs111447985 (10:105677911:C:A)




region 68: rs2732540 (11:35095486:A:G) and rs11032994 (11:35111157:G:A)




region 72: rs480958 (11:118577990:G:A) and rs4936441 (11:118725660:C:G)




region 73: rs11501246 (11:128314424:A:G) and rs684150 (11:128594161:C:T)




region 75: rs10845602 (12:12825237:A:G) and rs4251697 (12:12874462:G:A)




region 78: rs77465633 (12:111933545:C:A) and rs1131476 (12:113357209:G:A)




region 79: rs7139263 (12:118428976:G:A) and rs35743748 (12:121228629:C:CT)




region 84: rs7170151 (15:38846678:C:T) and rs12900640 (15:38974548:A:C)




region 94: rs447632 (16:85967640:A:G) and rs11117432 (16:86019271:G:A)




region 107: rs3794986 (19:52047314:G:T) and rs4801882 (19:52127053:G:A)



region 111: rs6641111 (X:12839790:C:G) and rs4830478 (X:12918288:G:A)




region 113: rs1059702 (X:153284192:A:G) and rs5945199 (X:153749857:G:A)



Supplementary Figure 3: Expression quantitative trait locus (eQTL) effect of rs66977652 on *SMG7* gene across a range of human tissues/cells in the GTEx v8 (<https://www.gtexportal.org>). Samples: the study sample size for each of the tissues. NES: normalized effect size. M-value is a posterior probability to quantify the association between rs66977652 and *SMG7* expression that was estimated in a Bayesian tool METASOFT (http://genetics.cs.ucla.edu/meta_jemdoc/). The rotated box plot shows the eQTL effect size and its 95% confidence interval in each tissue separately.



Supplementary Figure 4. Tissues (a) and cells (b) in which the expression of genes within SLE-associated loci are significantly enriched. Y-axis denotes the minus $\log_{10}$ -transformed enrichment P values. The tissues and cells significantly more expressing SLE-locus genes are colored in orange.



Supplementary Figure 5. Schematic diagram of luciferase expression vector constructs. To test the allele-specific enhancing activity of rs61759532, three constructs were used by adding no inserts (top; None) and three identical copies of the 24bp element around C alleles (middle; 3xC) or T alleles (bottom; 3xT) into the upstream of the minimal promoter (denoted as P) of the firefly luciferase gene of the luciferase vector (pGL4.26).

