

Online Supplements

Description of sample cohorts

Table S1. Descriptive statistics over subsamples.

Table S2. Correlations of ReMAP single mood item with BDI scores over subsamples.

Table S3. Correlations of ReMAP single sleep item with BDI sleep item over subsamples.

Description of sample cohorts

The sample for the current analyses were drawn from several ongoing longitudinal cohorts. Samples included stem from the Marburg/Münster Affective Disorder Cohort Study (MACS, n=47) (Vogelbacher et al., 2018), the Münster Neuroimage Cohort (MNC, n=17) (Dannlowski et al., 2016; Opel et al., 2019), two subsamples of the SFB-TRR58 cohort (n=81; Z02 Münster cohort and SpiderVR Münster cohort (Schwarzmeier et al., 2020)), and the TIP (n=28) cohort. All cohorts comprise healthy control (HC) participants, as well as different patient groups. Both, the MACS and the MNC cohorts include MDD and bipolar (BD) patients under current or former inpatient treatment. The SFB-TRR58 sample includes patients with a spider phobia (SP). The TIP sample includes patients with a social anxiety disorder (SAD), MDD, or with comorbid SAD and MDD.

References:

- Dannlowski, U., Kugel, H., Grotegerd, D., Redlich, R., Opel, N., Dohm, K., ... Baune, B. T. (2016). Disadvantage of Social Sensitivity: Interaction of Oxytocin Receptor Genotype and Child Maltreatment on Brain Structure. *Biological Psychiatry*, 80(5), 398–405. <https://doi.org/10.1016/j.biopsych.2015.12.010>
- Opel, N., Redlich, R., Dohm, K., Zaremba, D., Goltermann, J., Reppe, J., ... Dannlowski, U. (2019). Mediation of the influence of childhood maltreatment on depression relapse by cortical structure: a 2-year longitudinal observational study. *The Lancet Psychiatry*, 6(4), 318–326. [https://doi.org/10.1016/S2215-0366\(19\)30044-6](https://doi.org/10.1016/S2215-0366(19)30044-6)
- Schwarzmeier, H., Leehr, E. J., Böhnlein, J., Seeger, F. R., Roesmann, K., Gathmann, B., ... Dannlowski, U. (2020). Theranostic markers for personalized therapy of spider phobia: Methods of a bicentric external cross-validation machine learning approach. *International Journal of Methods in Psychiatric Research*, 29(2), e1812. <https://doi.org/10.1002/mpr.1812>
- Vogelbacher, C., Möbius, T. W. D., Sommer, J., Schuster, V., Dannlowski, U., Kircher, T., ... Bopp, M. H. A. (2018). The Marburg-Münster Affective Disorders Cohort Study (MACS): A quality assurance protocol for MR neuroimaging data. *NeuroImage*, 172, 450–460. <https://doi.org/10.1016/j.NEUROIMAGE.2018.01.079>

Table S1. Descriptive statistics over subsamples.

	Age		Sex	BDI _{ReMAP}		BDI _{stationary}		BDI difference (absolute)		Intervall (days)		Mood _{ReMAP}		Sleep _{ReMAP}	
	M (SD)	range		M (SD)	range	M (SD)	range	M (SD)	range	M (SD)	range	M (SD)	range	M (SD)	range
Full sample (n=173)	30.14	18-68	41/132	5.35	0-44	6.46	0-47	3.02	0-26	5.84	0.20-28.70	7.25	2-10	7.30	2-11
	(11.92)			(8.63)		(9.06)		(3.76)		(7.29)		(1.67)		(1.51)	
<i>BDI version subgroups</i>															
BDI-I _{stationary} (n=64)	37.17	20-68	14/50	7.20	0-44	7.48	0-43	3.25	0-26	8.07	0.20-28.70	6.91	2-10	7.22	2-10
	(14.28)			(10.54)		(10.18)		(4.61)		(8.07)		(1.87)		(1.68)	
BDI-II _{stationary} (n=109)	26.02	18-58	27/82	4.26	0-39	5.85	0-47	2.88	0-14	4.52	0.40-28.60	7.46	3-10	7.36	3-11
	(7.81)			(7.12)		(8.32)		(3.18)		(6.48)		(1.51)		(1.40)	
<i>Retest interval subgroups</i>															
<1 week interval (n=126)	28.79	18-58	30/96	4.59	0-44	6.18	0-47	2.77	0-22	1.82	0.20-7.80	7.33	2-10	7.26	2-10
	(10.72)			(8.16)		(9.08)		(3.35)		(1.91)		(1.64)		(1.50)	
>1 week interval (n=47)	33.77	18-68	11/36	7.38	0-37	7.19	0-41	3.68	0-26	16.59	8.40-28.70	7.04	3-10	7.43	4-11
	(14.17)			(9.57)		(9.08)		(4.66)		(5.15)		(1.75)		(1.53)	
<i>Disorder subgroups</i>															
healthy control (n=101)	28.92	18-68	26/75	1.89	0-17	2.97	0-13	2.25	0-13	5.14	0.20-28.70	7.67	4-10	7.36	3-10
	(11.52)			(3.12)		(3.30)		(2.55)		(6.83)		(1.31)		(1.27)	
affective disorders (n=57)	33.61	18-66	14/43	12.23	0-44	13.42	0-47	4.70	0-26	7.72	0.30-27.60	6.33	2-10	7.26	2-11
	(12.73)			(11.74)		(12.45)		(5.08)		(7.91)		(1.90)		(1.88)	
anxiety disorders (n=15)	25.20	18-43	1/14	2.47	0-8	3.47	0-13	1.80	0-8	3.33	0.40-22.30	8.07	5-10	7.07	4-10
	(8.06)			(2.47)		(4.00)		(2.57)		(6.68)		(1.44)		(1.49)	

Table S2. Correlations of ReMAP single mood item with BDI scores over subsamples.

	r	p	n
<i>All disorder groups</i>			
BDI-I _{ReMAP}	-.538	<.001	168
BDI-I _{stationary}	-.485	<.001	61
BDI-II _{stationary}	-.504	<.001	107
<i>Healthy controls</i>			
BDI-I _{ReMAP}	-.317	.001	98
BDI-I _{stationary}	-.170	.405	26
BDI-II _{stationary}	-.337	.004	72
<i>Affective disorders</i>			
BDI-I _{ReMAP}	-.494	<.001	58
BDI-I _{stationary}	-.462	.005	35
BDI-II _{stationary}	-.319	.138	23
<i>Anxiety disorders</i>			
BDI-I _{ReMAP}	-.092	.776	12
BDI-I _{stationary}	-	-	-
BDI-II _{stationary}	-.657	.020	12

Table S3. Correlations of ReMAP single sleep item with BDI sleep item over subsamples.

	r	p	n
<i>All disorder groups</i>			
BDI-I _{ReMAP} sleep item	-.310	<.001	166
BDI-I _{stationary} sleep item	-.279	.027	63
BDI-II _{stationary} sleep item	-.202	.042	102
<i>Healthy controls</i>			
BDI-I _{ReMAP} sleep item	-.387	<.001	97
BDI-I _{stationary} sleep item	-.179	.371	27
BDI-II _{stationary} sleep item	-.151	.211	70
<i>Affective disorders</i>			
BDI-I _{ReMAP} sleep item	-.395	.002	57
BDI-I _{stationary} sleep item	-.299	.077	36
BDI-II _{stationary} sleep item	-.379	.090	21
<i>Anxiety disorders</i>			
BDI-I _{ReMAP} sleep item	.275	.387	12
BDI-I _{stationary} sleep item	-	-	-
BDI-II _{stationary} sleep item	-.450	.165	11