

SUPPLEMENTARY INFORMATION FILES

Supplementary Text S1.

Calculating the Gini coefficient and bootstrap 95% confidence intervals using R version 3.6.3

- 1) Download and start R Version 3.6.3
- 2) Download **reldist** from the R library
- 3) Download **boot** from the R library
- 4) Command: **x <- c(n1,n2,n3,n4...)**
- 5) Command: **gini(x)** will give an output of the gini coefficient**
- 6) Command: **y <- boot(x,gini,500)** to set resampling of 500
- 7) Command: **quantile(y\$t, probs=c(0.025,0.975))**

Values of the Gini coefficient without the confidence intervals were also checked and confirmed using STATA version 15.3 and Microsoft Excel.

Supplementary Table S1. Gini coefficients of inequality among WHO world regions in the estimated burden (per 100,000 of the population) of major global diseases.

WHO World Region	Malaria 2018	HIV 2018	TB 2018	Hep C 2015	Cancer 2018	Diabetes 2014	Respiratory Deaths 2016	CVD Deaths 2016
Africa	23000	107	27	31	75.4	7100	18.1	112.8
Americas	670	16	11	6.4	373.2	8300	50.1	195.1
East Med	1000	7	1.3	62.5	97.4	13700	24.1	198.2
Europe	0.00	19	12	61.8	495.6	7300	48.9	435.6
SE Asia	490	9	3.5	14.8	100.7	8600	68.4	201.8
W Pacific	260	6.00	1.8	6.0	321.5	8400	62.2	290.0
Gini Coefficient (95% CI)	0.77 (0.66, 0.81)	0.56 (0.41, 0.58)	0.49 (0.37, 0.56)	0.43 (0.31, 0.49)	0.36 (0.26, 0.40)	0.12 (0.062, 0.15)	0.23 (0.14, 0.28)	0.22 (0.15, 0.26)

The Gini coefficients and 95% confidence intervals tabulated here are plotted in Figure 1. TB, tuberculosis; CVD, cardiovascular disease. Data are estimates as presented in WHO world reports and fact sheets, based on the most recent available, with references given in the paper.

Supplementary Table S2. No decline in the Gini coefficient of global malaria inequality based on estimated malaria cases as a percentage of Population at Risk between 2010 and 2018.

WHO World Region	2010	2011	2012	2013	2014	2015	2016	2017	2018
Africa	29.5	27.9	26.6	25.3	23.8	23.3	23.5	23.5	23.0
Americas	0.65	0.48	0.45	0.43	0.36	0.42	0.51	0.69	0.67
East Med	1.0	1.1	0.96	0.88	0.88	0.82	1.0	1.0	1.0
European	5x10 ⁻⁴	2x10 ⁻⁴	5x10 ⁻⁴	7x10 ⁻⁶	5x10 ⁻⁶	0.0	0.0	0.0	0.0
SE Asia	1.7	1.4	1.2	0.89	0.85	0.87	0.88	0.71	0.49
W Pacific	0.26	0.22	0.24	0.28	0.32	0.19	0.23	0.24	0.26
Gini	0.78	0.77	0.77	0.77	0.77	0.77	0.77	0.76	0.77
Coefficient (95% CI)	(0.67, 0.82)	(0.66, 0.81)	(0.66, 0.81)	(0.66, 0.81)	(0.66, 0.81)	(0.66, 0.81)	(0.66, 0.81)	(0.65, 0.80)	(0.66, 0.81)

Data are estimates of numbers of cases and populations at risk as presented in the WHO World Malaria Report 2019.

Supplementary Table S3. No decline in the Gini coefficient of malaria inequality among 16 West African countries based on estimated malaria cases as a percentage of Population at Risk between 2010 and 2018.

Country	2010	2011	2012	2013	2014	2015	2016	2017	2018
Benin	38.8	37.0	37.8	39.5	39.9	41.2	42.2	39.9	38.6
Burkina Faso	55.1	54.0	52.8	48.8	43.6	40.0	40.2	40.0	39.9
Cape Verde	0.04	0.01	0.0008	0.016	0.019	0.01	0.035	0.30	0.00
The Gambia	28.9	25.7	27.5	23.7	14.2	19.5	11.65	5.30	6.60
Ghana	36.4	38.0	37.43	34.93	31.58	27.7	23.60	21.3	22.4
Guinea	41.5	42.7	42.8	40.8	38.1	35.6	33.1	31.2	28.4
Guinea-Bissau	13.4	14.1	12.9	11.3	9.4	8.0	7.1	7.8	12.3
Ivory Coast	46.9	44.2	39.6	33.9	31.5	32.0	35.5	36.2	33.1
Liberia	34.6	33.0	30.8	31.7	33.8	34.7	38.6	40.1	36.2
Mali	38.4	40.5	43.6	45.3	44.1	39.2	38.4	38.7	38.7
Mauritania	3.9	4.8	2.8	3.3	4.9	6.2	7.2	5.5	3.9
Niger	42.6	42.8	43.1	42.0	40.0	37.0	35.9	35.7	35.7
Nigeria	39.9	37.3	34.8	32.9	31.4	29.6	28.1	28.3	29.2
Senegal	5.9	5.0	5.7	6.8	4.0	7.0	4.6	5.2	5.6
Sierra Leone	45.9	45.4	44.7	43.3	40.9	40.4	39.1	36.6	32.0
Togo	30.9	31.3	35.0	38.5	38.5	36.4	32.5	27.8	26.7
Gini Coefficient	0.28	0.28	0.28	0.28	0.29	0.27	0.30	0.32	0.31
(95% CI)	(0.19, 0.36)	(0.19, 0.36)	(0.17, 0.36)	(0.20, 0.38)	(0.18, 0.36)	(0.21, 0.37)	(0.21, 0.37)	(0.22, 0.36)	(0.22, 0.39)

The Gini coefficients and 95% confidence intervals tabulated here are plotted in Figure 2. Data are estimates of numbers of cases and populations at risk as presented in the WHO World Malaria Report 2019.

Supplementary Table S4. Gini coefficients show differences among West African countries in the levels of sub-national variation in malaria parasite prevalence in community surveys of children <5 years of age


Prevalence in the major administrative units surveyed within each of the countries:							
Nigeria		Ghana		Burkina Faso		Sierra Leone	
Benue	44.5	Western	23.5	Boucle	23.3	Kailahun	45
Kogi	5.4	Central	30.2	Cascades	13	Kenema	37.7
Kwara	26.4	Accra	4.8	Centre	7.2	Kono	37.5
Nasarawa	35.9	Volta	27.5	E Centre	12.4	Bombali	37.6
Niger	33.5	Eastern	31.3	N Centre	18.2	Kambia	48.3
Plateau	35.8	Ashanti	16.6	W Centre	25.4	Koinadugu	57.9
Adamawa	34.7	Brong Ahafo	22.4	S Centre	12.3	Port Loko	58.5
Bauchi	19.6	Northern	25.2	East	17.6	Tonkolili	55.7
Gombe	28.6	Upper East	14.7	Hauts-Bas	12.7	Bo	39.7
Taraba	42.9	Upper West	21.5	North	11.4	Bonthe	26.1
Yobe	18.9			Plateau C	8.4	Moyamba	39.9
Jigawa	27.9			Sahel	18.2	Pujehun	46.8
Kaduna	36.7			South-West	38.9	W Rural	34.9
Oyo	19.2					W Urban	6.3
Abia	8.2						
Anambra	10.2						
Ebonyi	30						
Enugu	10.5						
Imo	5.1						
Akwa Ibom	22.8						
Bayelsa	31.4						
Cross River	26.1						
Delta	20.4						
Edo	18.6						
Rivers	7.3						
Ekiti	28.8						
Lagos	0.0						
Osun	33.4						
Kano	27.7						
Katsina	27.8						
Kebbi	63.6						
Sokoto	46.6						
Zamfara	62.6						
Ogun	14.7						
Ondo	21.3						
Gini	0.30		0.19		0.25		0.17
(95% CI)	(0.26, 0.35)		(0.12, 0.25)		(0.19, 0.29)		(0.12,0.22)

The Gini coefficients and 95% confidence intervals tabulated here are plotted in Figure 3. Data are from national Multiple Indicator Surveys with references in the paper.

Supplementary Table S5. Gini coefficients show extent of variation in community prevalence of *P. falciparum* among 16 villages in the Garki Project in northern Nigeria in eight cross-sectional surveys during the pre-intervention phase of the project.

Village	Percent prevalence at each of eight surveys							
	1	2	3	4	5	6	7	8
Mukawa	54.3	48.0	27.4	39.4	36.6	37.8	39.2	38.0
Matsari	63.6	41.3	34.6	39.3	41.9	46.5	30.0	35.6
Rafin Marke	57.8	32.2	44.1	43.6	58.4	47.6	33.6	37.1
Kukar	66.6	40.0	36.2	40.3	57.7	45.4	30.2	41.9
UngwarBako	64.1	39.3	35.5	40.8	49.0	50.2	37.0	37.6
Kargo Kudu	50.1	24.7	32.0	42.1	47.6	41.0	23.5	30.3
Nasakar	62.5	38.8	42.4	43.7	60.5	51.2	20.7	37.1
Jaya	57.3	40.0	22.1	33.4	39.1	46.1	32.1	21.3
Sugungun	54.4	47.0	41.2	42.9	53.3	55.6	47.6	44.8
Karama	55.8	49.6	43.0	38.6	55.1	55.5	43.2	39.7
Batakashi	60.2	44.2	50.0	49.0	50.0	67.1	39.6	43.2
Nabanawa	63.2	41.0	40.8	44.0	58.6	46.5	34.1	38.4
Ajura	49.7	33.5	37.6	38.9	56.7	44.2	32.4	34.7
Barebari	60.8	50.6	47.6	46	53.4	61.5	42.3	39.5
Kwaru	55.6	39.6	46.4	45.9	54.1	63.4	49.5	38.5
Tafin Sale	45.7	26.9	31.4	35.3	40.0	39.8	35.5	37.3
Gini Coefficient	0.056	0.10	0.11	0.053	0.082	0.093	0.17	0.070
(95% CI)	(0.043, 0.062)	(0.075, 0.12)	(0.088, 0.13)	(0.042, 0.058)	(0.059, 0.09)	(0.077, 0.11)	(0.11, 0.20)	(0.044, 0.09)

The Gini coefficients and data tabulated here are plotted in Figure 4. Each of the surveys was approximately 10 weeks apart, and variation among the villages was highest in the dry season.



Supplementary Figure S1.

High Gini coefficients of variation in *P. falciparum* prevalence among villages in The Gambia with seasonal malaria transmission. Variation was analysed among 20 villages each surveyed at two seasonal times. There were 3860 individuals sampled in the wet season and 3707 in the dry season (prevalence measurements in each of the villages are previously published). The coastal area had 6 villages sampled, while the mid country area and east country areas each had 7 villages sampled. The east country area showed statistically greater variation among villages in the dry season.